

1                   FOOD AND DRUG ADMINISTRATION  
2                   CENTER FOR DRUG EVALUATION AND RESEARCH  
3                   ARTHRITIS ADVISORY COMMITTEE  
4  
5  
6  
7                   WEDNESDAY, SEPTEMBER 16, 2009  
8                   8:30 a.m. to 3:00 p.m.  
9  
10  
11                  Holiday Inn Gaithersburg  
12                  Two Montgomery Village Avenue  
13                  Gaithersburg, Maryland  
14  
15  
16  
17  
18  
19  
20  
21  
22

1 Diane Aronson (Consumer Representative)

2 Consumer Advocacy

3 P.O. Box 410305

4 Cambridge, MA 02141

5

6 Lenore Buckley, M.D., M.P.H.

7 Professor of Internal Medicine and Pediatrics

8 Virginia Commonwealth University

9 School of Medicine

10 P.O. Box 980102

11 Richmond, VA 23298

12

13 Nancy Olsen, M.D.

14 Professor of Internal Medicine

15 McGee Foundation Chair in Arthritis Research

16 University of Texas Southwestern Medical School

17 5323 Harry Hines Boulevard

18 Dallas, TX 75390

19

20

21

22

1 Kathleen O'Neil, M.D. (Chair)

2 Associate Professor of Pediatrics

3 Department of Pediatrics

4 Division of Rheumatology

5 University of Oklahoma College of Medicine

6 940 N.E. 13th Street, Bielstein 3B3303

7 Oklahoma City, OK 73104

8

9 Kenneth Saag, M.D.

10 Associate Professor

11 Department of Medicine

12 Division of Clinical Immunology and Rheumatology

13 The University of Alabama at Birmingham

14 820 Faculty Office Tower

15 510 20th Street South

16 Birmingham, AL 35294

17

18 **TEMPORARY VOTING MEMBERS**

19 William Brackney (Patient Representative)

20 Henderson, NV 89011

1 Mustafa Haque, M.D.

2 Clinical Assistant Professor

3 Department of Orthopedic Surgery

4 Georgetown University Hospital

5 Hand Surgeon

6 Summit Orthopaedic Surgery

7 5530 Wisconsin Avenue, Suite 1660

8 Chevy Chase, MD 20815

9

10 Saul Kaplan, M.D.

11 Hand Surgeon

12 Mid-Atlantic Permanente Medical Group

13 6501 Loisdale Court

14 Springfield, VA 22150

15

16 Kathleen Mazor, Ed.D.

17 Associate Professor of Medicine

18 University of Massachusetts Medical School

19 630 Plantation Street

20 Worcester, MA 01605

21

22

1 Timothy McAlindon, M.D., M.P.H, M.R.C.P.

2 Professor of Medicine and Natalie V. Zucker and

3 Milton O. Zucker Chair of Rheumatology

4 Tufts University School of Medicine

5 Chief, Division of Rheumatology

6 Tufts Medical Center

7 35 Kneeland Street 4th Floor

8 Boston, MA 02111

9

10 William Swartz, M.D., F.A.C.S.

11 Clinical Associate Professor of Surgery

12 University of Pittsburgh, Department of Plastic

13 Surgery

14 Specialties of Plastic, Hand and Microsurgery, P.C.

15 5750 Centre Ave. Suite 180

16 Pittsburgh, PA 15206

17

18

19

20

21

22

1    Michael Weisman, M.D.

2    Division of Rheumatology

3    Department of Medicine

4    Cedars-Sinai Medical Center

5    8700 Beverly Boulevard

6    Los Angeles, CA 90048

7

8    **FDA Non-Voting**

9    Curtis Rosebraugh, M.D., M.P.H

10   Director, Office of Drug Evaluation II

11   CDER/FDA

12

13   Bob Rappaport, M.D.

14   Director, Division of Anesthesia, Analgesia and

15   Rheumatology Products

16   CDER/FDA

17

18   Sarah Okada, M.D.

19   Clinical Team Leader, Division of Anesthesia,

20   Analgesia and

21   Rheumatology Products

22   CDER/FDA

1 Eric Brodsky, M.D.

2 Clinical Reviewer, Division of Anesthesia, Analgesia

3 and Rheumatology Products

4 CDER/FDA

5

6 Kathryn O'Connell, M.D., Ph.D.

7 Drug Risk Management Analyst

8 OSE/CDER/FDA

9

10

11

12

13

14

15

16

17

18

19

20

21

22

1 I N D E X

2	AGENDA ITEM	PAGE
3	Call to Order and Introduction	
4	Kathleen O'Neil, M.D.	9
5	Conflict of Interest Statement	
6	Nicole Vesely, Pharm.D.	12
7	Opening Remarks	
8	Sarah Okada, M.D	18
9	Sponsor Presentation	
10	Benjamin Del Tito, Ph.D.	20
11	F. Thomas Kaplan, M.D.	25
12	Anthony DelConte, M.D.	39
13	James Tursi, M.D.	48
14	Anthony DelConte, M.D.	75
15	Questions to the Sponsor	77
16	FDA Presentation	
17	Eric Brodsky, M.D.	113
18	Kathryn O'Connell, M.D., Ph.D.	130
19	Questions to the FDA	136
20	Open Public Hearing	165
21	Questions to Committee and Discussion	214
22	Adjourn	269



1                   P R O C E E D I N G S

2                   8:30 a.m.

3                   DR. O'NEIL: Good morning. My name is  
4 Kathleen O'Neil, and I'm an associate professor of  
5 pediatrics and rheumatology at the University of  
6 Oklahoma in Oklahoma City. We are ready to begin the  
7 meeting of the FDA Arthritis Advisory Committee.

8                   I would like to start by asking everyone at  
9 the table to introduce themselves, starting here with  
10 Dr. Haque.

11                  DR. HAQUE: My name is Mustafa Haque. I'm a  
12 practicing orthopedic hand and upper extremities  
13 surgeon in Chevy Chase, Maryland.

14                  DR. SWARTZ: Good morning. I'm Bill Swartz  
15 from Pittsburgh, Pennsylvania. I'm a practicing hand  
16 surgeon, been in practice for 30 years.

17                  DR. S. KAPLAN: Saul Kaplan, Fairfax,  
18 Virginia. I'm an orthopedic hand surgeon in practice.

19                  DR. MAZOR: Kathy Mazor. I'm associate  
20 professor at the University of Massachusetts Medical  
21 School. I'm not a physician. My background's in  
22 education, psychometrics, patient education and

1 physician-patient communication.

2 DR. McALINDON: I'm Tim McAlindon. I'm  
3 chief of rheumatology at Tufts Medical Center, and  
4 professor of medicine at Tufts University School of  
5 Medicine. I'm a clinical rheumatologist. I also do  
6 clinical investigations into rheumatic diseases.

7 DR. OLSEN: I'm Nancy Olsen. I'm a  
8 professor of medicine at the University of Texas  
9 Southwestern Medical School in Dallas. I'm a  
10 rheumatologist in academic practice, and I'm  
11 interested in autoimmune diseases.

12 DR. BUCKLEY: I'm Lenore Buckley. I'm a  
13 professor of medicine and pediatric at Virginia  
14 Commonwealth University, and I do both adult and  
15 pediatric rheumatology.

16 DR. VESELY: Nicole Vesely, designated  
17 federal official, Arthritis Advisory Committee.

18 DR. SAAG: Good morning. I'm Ken Saag. I'm  
19 a professor of medicine and epidemiology at the  
20 University of Alabama at Birmingham, where I direct  
21 the AHRQ-funded Center for Education and Research in  
22 Therapeutics.

1 MS. ARONSON: I'm Diane Aronson. I'm a  
2 consumer representative, standing member of the  
3 Arthritis Committee. I'm from Cambridge,  
4 Massachusetts.

5 MR. BRACKNEY: Bill Brackney. I'm from  
6 Henderson, Nevada, and I'm a patient representative.

7 DR. WEISMAN: I'm Michael Weisman, director  
8 of the Division of Rheumatology at Cedars-Sinai  
9 Medical Center, and professor of medicine at UCLA  
10 School of Medicine. And I'm a rheumatologist,  
11 interested in outcomes and risk for rheumatic  
12 diseases.

13 DR. O'CONNELL: Good morning. My name is  
14 Kathryn O'Connell. I'm a medical officer in FDA's  
15 Division of Risk Management.

16 DR. BRODSKY: Good morning. My name is Eric  
17 Brodsky. I'm a medical officer in rheumatology at the  
18 FDA.

19 DR. OKADA: Hi, Sarah Okada, clinical team  
20 leader of the Division of Anesthesia, Analgesia and  
21 Rheumatology Products, and I'm an adult  
22 rheumatologist.

1 DR. RAPPAPORT: Good morning. I'm Bob  
2 Rappaport. I'm the director of that division.

3 DR. ROSEBRAUGH: Curt Rosebraugh, director  
4 of the Office of Drug Evaluation II.

5 DR. VESELY: For topics such as those being  
6 discussed at today's meeting, there are often a  
7 variety of opinions, some of which are quite strongly  
8 held. Our goal is that today's meeting will be a fair  
9 and open forum for discussion of these issues, and  
10 that individuals can express their views without  
11 interruption. Thus, as a gentle reminder, individuals  
12 will be allowed to speak into the record only if  
13 recognized by the Chair. We look forward to a  
14 productive meeting.

15 In the spirit of the Federal Advisory  
16 Committee Act and the Government in the Sunshine Act,  
17 we ask that the Advisory Committee members take care  
18 that their conversations about the topic at hand take  
19 place in the open forum of the meeting. We are aware  
20 that members of the media are anxious to speak with  
21 the FDA about these proceedings. However, FDA will  
22 refrain from discussing the details of this meeting

1 with the media until its conclusion. Also, the  
2 Committee is reminded to please refrain from  
3 discussing the meeting topic during breaks or lunch.  
4 Thank you.

5 And now for the conflict of interest  
6 statement. The Food and Drug Administration is  
7 convening today's meeting of the Arthritis Drugs  
8 Advisory Committee under the authority of the Federal  
9 Advisory Committee Act of 1972. With the exception of  
10 the industry representative, all members and temporary  
11 voting members of the Committee are special government  
12 employees or regular federal employees from other  
13 agencies, and are subject to federal conflict of  
14 interest laws and regulations.

15 The following information on the status of  
16 this Committee's compliance with federal ethics and  
17 conflict of interest laws covered by but not limited  
18 to those found at 18 USC Section 208 and Section 712  
19 of the Federal Food, Drug and Cosmetic Act is being  
20 provided to participants in today's meeting and to the  
21 public.

22 The FDA has determined that members and

1 temporary voting members of this Committee are in  
2 compliance with federal ethics and conflict of  
3 interest laws. Under 18 USC Section 208, Congress has  
4 authorized FDA to grant waivers to special government  
5 employees and regular federal employees who have  
6 potential financial conflicts, when it is determined  
7 that the agency's need for particular individual  
8 services outweighs his or her potential financial  
9 conflict of interest.

10 Under Section 712 of the FD&C Act, Congress  
11 has authorized FDA to grant waivers to special  
12 government employees and regular federal employees  
13 with potential financial conflicts when necessary to  
14 afford the Committee essential expertise.

15 Related to the discussion of today's  
16 meeting, members and temporary voting members of this  
17 Committee have been screened for potential financial  
18 conflicts of interest of their own as well as those  
19 imputed to them, including those of their spouses or  
20 minor children, and for purposes of 18 USC Section  
21 208, their employers.

22 These interests may include investments,

1 consulting, expert witness testimony, contracts,  
2 grants, CRADAs, teaching, speaking, writing, patents  
3 and royalties and primary employment.

4 Today's agenda involves discussion of  
5 collagenase clostridium histolyticum for the proposed  
6 treatment of advanced Dupuytren's disease under  
7 Biologics License Application 125338, sponsored by  
8 Auxilium Pharmaceuticals. This topic is a particular  
9 matter involving specific parties. Based on the  
10 agenda for today's meeting and all financial interests  
11 reported by the Committee members and temporary voting  
12 members, no conflict of interest waivers have been  
13 issued in connection with this meeting.

14 To ensure transparency, we encourage all  
15 standing members and temporary voting members to  
16 disclose any public statements that they have made  
17 concerning the product at issue. We would like to  
18 remind members and temporary voting members that if  
19 the discussions involve any products or firm not  
20 already on the agenda for which an FDA participant has  
21 a personal or imputed financial interest, the  
22 participants need to exclude themselves from such

1 involvement, and their exclusion will be noted for the  
2 record.

3 The FDA encourages all other participants to  
4 advise the Committee of any financial relationships  
5 that they may have with any firms at issue. We also  
6 just wanted to note that there is not an industry  
7 representative for this meeting.

8 Thank you.

9 DR. O'NEIL: Our first speaker this morning  
10 will be Dr. Bob Rappaport, the director of the  
11 Division of Anesthesia, Analgesia and Rheumatology  
12 Products at CDER FDA.

13 DR. RAPPAPORT: Thank you. Good morning,  
14 everybody. I don't think a lot of people in the  
15 public realize the time and effort and resources that  
16 the people who sit on our committees give us, give the  
17 FDA and the American public by participating. Our  
18 Committee members serve for periods from anywhere from  
19 two to four years, and during that time, they may  
20 cover numerous meetings and help us in other projects.  
21 And it's really -- as they're well-aware, they're paid  
22 a pittance for doing this. And we really do



1 appreciate their service.

2           So on behalf of the FDA, I'd like to take a  
3 brief moment to recognize one of our committee members  
4 whose term expires at the end of September.

5           Dr. Saag, would you come up?

6           Dr. Ken Saag has served on the Arthritis  
7 Advisory Committee since August of 2006. He's a  
8 professor in the Department of Medicine, Division of  
9 Clinical Immunology and Rheumatology at the University  
10 of Alabama at Birmingham. His expertise in rheumatoid  
11 arthritis and osteoporosis has brought a valuable  
12 knowledge base to this Committee's discussions, and  
13 his experience as both a clinician and researcher has  
14 proved invaluable.

15           So in appreciation of this service, the FDA  
16 would like to recognize Dr. Saag's service with this  
17 plaque.

18           Thank you very much.

19           DR. SAAG: Thank you very much, Bob.

20           DR. O'NEIL: Thank you, Dr. Rappaport.

21           We will begin the business portion of the  
22 meeting with some opening remarks from Dr. Sarah

1 Okada, who is the clinical team leader, also at the  
2 Division of Anesthesia, Analgesia and Rheumatology  
3 Products.

4 DR. OKADA: Good morning, everyone. I'd  
5 like to welcome you all and thank our Advisory Panel  
6 once again for taking time out of your busy schedules  
7 to join us for today's meeting. The topic for  
8 discussion today is Auxilium Pharmaceutical's  
9 clostridial collagenase, also known as AA4500 or  
10 Xiaflex, as a nonsurgical treatment of Dupuytren's  
11 contractures.

12 So in addition to our esteemed Arthritis  
13 Advisory Committee regulars, whose continued support  
14 we greatly appreciate, we have several special guests  
15 joining our panel today. Since the management of  
16 Dupuytren's disease has historically been the purview  
17 of our surgical colleagues, we're fortunate today to  
18 have several hand surgeons on the panel. So I'd like  
19 to extend a special thanks to Drs. Haque, Swartz and  
20 Kaplan for joining us today.

21 We're also fortunate to have two members of  
22 the Drug Safety and Risk Management Advisory

1 Committee, Dr. McAlindon and Dr. Mazor.

2           Today, we'll be starting off with the  
3 sponsor presentations, which will cover in depth the  
4 efficacy and safety data for AA4500, in addition to  
5 their proposed risk management activities. Then the  
6 main FDA presentation by Dr. Eric Brodsky will contain  
7 only a brief discussion of efficacy, which is not in  
8 question, and focus more on the safety data in the  
9 trials and concerns relative to generalizability of  
10 study results.

11           Following this, we will have a brief  
12 overview of risk management considerations in the FDA  
13 approval process by Dr. Kathryn O'Connell from the  
14 Office of Surveillance and Epidemiology.

15           Finally, we will be asking the Committee to  
16 discuss the proposed training for health care  
17 professionals in clinical practice, whether this  
18 training is adequate, and what factors will facilitate  
19 the assimilation of the training for the safe and  
20 effective use of this product, as well as to discuss  
21 the overall risk/benefit profile of the product and  
22 whether you recommend the product be approved.

1           Again, our deepest thanks to the panel, and  
2   we look forward to hearing your views. So without  
3   further adieu, I'll turn this back over to Dr. O'Neil.

4           DR. O'NEIL: Thank you. We will now move on  
5   to our presentation by Auxilium Pharmaceuticals,  
6   sponsors of this product. The first speaker who will  
7   introduce the product is Dr. Benjamin Del Tito, senior  
8   vice president, Quality and Regulatory Affairs, at  
9   Auxilium Pharmaceuticals.

10          DR. DEL TITO: Good morning. My name is  
11   Ben Del Tito, and I am the senior vice president of  
12   Quality and Regulatory Affairs for Auxilium. I would  
13   like to thank the Arthritis Advisory Committee and the  
14   FDA on behalf of Auxilium Pharmaceuticals for  
15   providing us with the opportunity to discuss our  
16   complete drug development program with you; that being  
17   AA4500, collagenase clostridium histolyticum.

18          After my brief introduction, I will turn it  
19   over to Dr. Tom Kaplan, who is an orthopedic hand  
20   surgeon and one of our Phase 3 clinical investigators,  
21   and he will discuss the disease state as well as its  
22   current management.

1           That will be followed by Dr. Tony DelConte,  
2   who is Auxilium's chief medical officer, and he will  
3   discuss AA4500 clinical efficacy.

4           He'll turn it over to Dr. Jim Tursi, who is  
5   Auxilium's vice president of Clinical Affairs, and he  
6   will discuss AA4500 clinical safety and our risk  
7   management activities.

8           And finally, Dr. DelConte will return to the  
9   podium to discuss the overall summary of our program.

10          We use our hands constantly for various  
11   tasks in our daily lives from the moment we wake up in  
12   the morning until we go to bed at night. Activities  
13   such as writing with a pen, typing on a computer or a  
14   BlackBerry -- those afflicted with the debilitating  
15   disease known as Dupuytren's struggle every day with  
16   these same tasks that we take for granted. I'd like  
17   to ask the panel to please keep that in mind as we  
18   discuss our program with you today.

19          We're here to discuss AA4500, collagenase  
20   clostridium histolyticum for injection. The proposed  
21   indication is the treatment of advanced Dupuytren's  
22   disease, and that can be defined as a progressive

1 disease resulting in a fixed flexion deformity or a  
2 contracture in one of several joints, most commonly  
3 the last two digits of the hand. A Dupuytren's cord  
4 is an abnormal collagen deposition in the palm of the  
5 hand, and it results in the contracture.

6           Now, the current treatment for Dupuytren's  
7 disease is surgery. What we would like to discuss  
8 today with you is an alternative to surgery, a novel  
9 option for physicians who treat Dupuytren's disease.  
10 AA4500 is a new molecular entity, and it's a first in  
11 class biological.

12           AA4500 consists of two collagenases. These  
13 are enzymes that are mixed in a fixed ratio. It's a  
14 naturally produced product by the bacteria -- the gram  
15 positive bacterium known as clostridium histolyticum,  
16 and the two enzymes are referred to as AUX-I and  
17 AUX-II, and these cleave the collagen substrate.  
18 Clostridium collagenases act in a complementary  
19 manner.

20           AA4500 dosage form is presented in a sterile  
21 lyophilized powder in single-use vials, and it's  
22 accompanied by a second vial which contains a sterile

1     diluent, consisting of calcium chloride and sodium  
2     chloride. The calcium is a required cofactor for  
3     enzymatic activity.

4             A single dose consists of 0.58 milligrams  
5     from a single-use vial. Now, this is injected  
6     directly into the cord, or intralesionally, and it's  
7     followed by a finger extension or manipulation after  
8     24 hours to disrupt the cord. Each cord can receive  
9     one injection at four-week intervals, for up to a  
10    total of three injections.

11            As I mentioned, AA4500 is administered by  
12    direct injection into the Dupuytren's cord. This  
13    consists of a third of the dose injected three times  
14    with close proximity into the cord. Once injected,  
15    AA4500 acts locally.

16            This illustrates the complementary activity  
17    of AA4500 components. Starting with the left panel,  
18    we see the action of AUX-I, which is a Class 1  
19    collagenase, exhibiting activity against intact  
20    collagen, cleaving at the ends of the collagen  
21    molecule shown here -- this being the amino terminus  
22    of the collagen and the carboxy terminus of the

1 collagen.

2           Then, AUX-II in the middle panel is a Class  
3 2 collagenase, and this exhibits activity against  
4 collagen peptides or fragments of collagen, and  
5 cleaves internally on the collagen molecule.  
6 Combining these two enzymes to form AA4500 results in  
7 a more complete degradation, because cleavage occurs  
8 on multiple sites on the collagen molecule.

9           A few regulatory achievements are shown on  
10 this slide. The Investigational New Drug application,  
11 or IND, was filed in 1994. An agreement with the  
12 agency was made on the dose selected, the .058  
13 milligrams equivalent to 10,000 units during the end  
14 of Phase 2 meeting in 2001. Auxilium licensed the  
15 product in 2004 with a subsequent IND transfer. Our  
16 Biologics License Application, or BLA, was filed in  
17 February of 2009, and it was accepted for filing by  
18 the FDA with a priority designation in April of 2009.

19           Joining us in the sponsors panel, we have a  
20 few outside experts: Dr. Tom Kaplan, who I mentioned  
21 earlier as an orthopedic surgeon, hand surgeon from  
22 Indiana University School of Medicine. We also have



1 Dr. Philip Waller, who is a practicing rheumatologist  
2 from Houston, Texas, and we also have Mr. Paul  
3 Chamberlain, an expert immunologist from the NDA  
4 Regulatory Sciences group in the UK.

5           Joining us from Auxilium in addition to  
6 Drs. Tursi, DelConte and myself are Dr. Ted Smith, who  
7 is Auxilium's vice president of Biometrics, and  
8 Dr. Susan Emeigh Hart, Auxilium's senior director of  
9 Drug Safety and Metabolism.

10           And with that, I would like to turn it over  
11 to Dr. Tom Kaplan, who will discuss the disease state  
12 and its current management. Dr. Kaplan.

13           DR. T. KAPLAN: Thank you, Dr. Del Tito.

14           Before we discuss the data on AA4500, I'd  
15 like to take a few minutes with the Committee to  
16 review what Dupuytren's disease is, and what our  
17 current treatment methods for it are. Dupuytren's  
18 disease is a progressive, fibroproliferative disorder  
19 that affects the tissue in the palm of our hands and  
20 fingers. As it develops, cords and nodules form in  
21 the hand contracting the fingers, drawing them down  
22 towards the palm. We typically see it most commonly

1 in the ring and small fingers initially, and see it in  
2 patients approximately 50 percent of the time  
3 bilaterally.

4           The pathoanatomy of Dupuytren's disease is  
5 that it affects the palmar fascia. This fascia is a  
6 layer of tissue underneath the skin that extends up  
7 the palm and into the fingers. It's above the flexor  
8 tendons and neurovascular bundles in the palm, and as  
9 that tissue in the palm is organized into a triangular  
10 configuration, there are these longitudinal bands that  
11 run along the palmar fascia. It is these bands that  
12 become diseased with progressive collagen deposition,  
13 and cords will form in these bands.

14           As these bands extend up towards the finger,  
15 they become somewhat more complex. They'll bifurcate  
16 or trifurcate actually and go towards the web of the  
17 digit. They can extend along the sides of the digit,  
18 and at this level, will actually wrap around the  
19 nerves and arteries as they go up into the finger.

20           In the early stages of the disorder,  
21 fibroblasts begin proliferating and differentiate into  
22 the pathognomonic cell of Dupuytren's disease called

1 the myofibroblast. These myofibroblasts actually have  
2 smooth muscle components and have a contractile  
3 ability. And this is when we first see nodule  
4 formation, and typically in the palm.

5 In the intermediate phase, the  
6 myofibroblasts begin to align along lines of tension  
7 in the palm of the hand, and with progressive collagen  
8 deposition, cords begin to form. And in the advanced  
9 disease, these cords begin to shorten, causing a  
10 finger contracture.

11 Nodules seen here in the palm of the hand  
12 most typically are seen over the MP joint in the palm.  
13 They are usually painless, but many patients will  
14 present at the nodule stage not sure of what is  
15 growing in their hand, concerned that it may be  
16 something more serious like a tumor or a cancer  
17 condition. These usually do not bother patients other  
18 than their appearance, but some patients will have  
19 pain associated with the nodules, especially if the  
20 nodules are particularly active, or patients who have  
21 to do a lot of repetitive gripping activities.

22 Other times, also less common, the nodules

1 will actually cause an irritation of the underlying  
2 flexor tendons, causing a flexor tenosynovitis.

3           Following nodule formation, we typically see  
4 the progression into cords, and you can see these  
5 well-defined bands extending up the digits. These  
6 cords, as they travel up the palm and into the finger,  
7 connect to the skin, and we'll oftentimes see the skin  
8 draw down into the hand, forming a pit. They'll go  
9 across down towards the joint, causing contracture,  
10 and they can course around the nerves and arteries.

11           As these cords shortened, contractures form.  
12 The cords in the palm typically cause contractures of  
13 the MP joint, as seen in this patient, where the cord  
14 comes down the center of the finger, the big nodule  
15 here and draws this joint down into a contracted  
16 position. The cords in the finger oftentimes cause  
17 contractures of the proximal and distal  
18 interphalangeal joints.

19           The prevalence of the disease varies  
20 depending on the population that we're looking at.  
21 It's more common in the northern -- in the Caucasian  
22 population, particularly patients of Northern European

1   ancestry. It is a disease of adult life, and affects  
2   men much more frequently than women.

3               The etiology is not completely understood.  
4   There have been many associations with Dupuytren's  
5   disease, and it's felt most likely to represent a  
6   genetic condition which exhibits an autosomal dominant  
7   pattern with variable penetrance. Familial clustering  
8   is seen.

9               Other associations which have been reported  
10   in the literature include those that cause tissue  
11   ischemia such as smoking and diabetes; trauma,  
12   especially manual laborers who may have repetitive  
13   microtears of that palmar fascia; epilepsy, where the  
14   drugs used to treat epilepsy such as Dilantin and  
15   alcoholism.

16              So what is the impact of Dupuytren's disease  
17   on the patient? Well, we know what science and the  
18   literature tell us. This is a Sollerman test, which  
19   looks at many common day-to-day activities such as  
20   putting a key in a lock and turning it, picking up a  
21   coin from a table, unscrewing the lid of a jar and  
22   buttoning your shirt. What Sinha looked at in 2002

1 was to correlate the Sollerman scale or test results,  
2 the maximum score of which was 80, and what they found  
3 was in patients with a less severe contracture, scores  
4 tended to be higher. And as the contracture  
5 progressed and worsened, their scores tended to  
6 deteriorate.

7           They also correlated this with a  
8 postoperative function, and found that the scores  
9 would increase again after surgery to correct the  
10 deformity.

11           The patients do the best job of telling how  
12 Dupuytren's disease affects them. Most commonly,  
13 patients describe difficulties doing their daily  
14 activities, particularly with personal hygiene, such  
15 as washing their face, combing their hair, tying a tie  
16 and shaking hands. It can also affect patients' jobs,  
17 particularly patients who have to get their hands in  
18 tight spaces, who have to wear gloves for their job or  
19 use a keyboard.

20           And because this is a disease of advanced  
21 years, many patients are retired and looking towards  
22 their hobbies in their retirement, and can no longer

1 do the sports that they were looking forward to or  
2 enjoying, or hobbies such as woodworking or playing a  
3 musical instruments.

4           Because there's no cure for Dupuytren's  
5 disease, treatment is based upon the severity of the  
6 disorder in the patient. Until a patient has a  
7 functional limitation of their hand, we usually  
8 recommend observation. This can be -- if someone has  
9 a painful nodule, oftentimes a massage may be helpful.  
10 Occasionally, corticosteroids are used for a painful  
11 nodule as well.

12           But we reserve treatment of the contracture  
13 until the contracture is bad enough, because there's  
14 limitations with all of our treatments. A quick test  
15 is when a patient can't get their hand flat on a table  
16 anymore, we typically think that their contracture has  
17 advanced to the point that intervention is warranted.  
18 A rough scale, that's an MP contracture of  
19 approximately 30 degrees, or a PIP joint contracture  
20 of approximately 20 degrees.

21           The current treatment options of surgery are  
22 either a fasciotomy, which involves division of the

1 cord at one or more locations; fasciectomy, which  
2 involves excision of the entire diseased cord which is  
3 causing the contracture; or a dermofasciectomy, which  
4 involves excision of that cord and the overlying skin  
5 with it, which then necessitates placement of a skin  
6 graft on top of the defect.

7           This last option is typically reserved for  
8 patients who have surgery previously and have had  
9 recurrent disease.

10           Fasciotomy can be performed either in an  
11 open end or percutaneously. This is a patient who had  
12 a contracture of his small finger at both the MP and  
13 PIP joints and was not willing to undergo the rigors  
14 of a more formal surgical procedure or the  
15 postoperative recuperation necessary. So through  
16 three small incisions along the palm and into the  
17 finger, we sectioned the cord, and was able to obtain  
18 this type of correction we got the MP joint fairly  
19 well-corrected. However, you can note there's still  
20 some mild contracture left at the PIP joint.

21           The problem with a fasciotomy is that  
22 recurrence is very frequent. In a study in 1997,



1 Duthie, et al, looked at 82 patients with an average  
2 preoperative contracture of 71 degrees. At ten-year  
3 follow-up, one-third of the patients had no further  
4 treatment, and their contracture had worsened to the  
5 point of 57 degrees. Most interesting is that  
6 two-thirds of the patients required further treatment  
7 at an average of five years after the index procedure.  
8 And by that time, their contracture had worsened to 85  
9 degrees.

10           This is also being performed more commonly  
11 with a needle procedure. The advantage of this is a  
12 quicker recuperation, less morbidity associated with  
13 the procedure, and you use a small needle in order to  
14 section that corridor at one or more locations.  
15 Unfortunately, although it's more tolerable to  
16 patients, it still has the problems of high recurrence  
17 rate.

18           In three various studies from 1993 to 2006,  
19 recurrence rates varied from 50 to 65 percent at  
20 average of three years. It's also associated with  
21 numerous potential complications, as it is a  
22 relatively blind procedure. Nerves can be sectioned

1 as well as arteries. Skin fissuring has been reported  
2 as well as flexor tendon injury.

3           So our current mainstay of treatment in the  
4 U.S. is subtotal palmar fasciectomy; that is, excision  
5 of the diseased cord which is causing the contracture.  
6 You see this is a typical patient preoperatively.  
7 He's asked to open and close your hand, and you see  
8 the limitation of both the MP and PIP joint levels.  
9 Again, when we think about surgery, when the MP joint  
10 can't extend to more than 30 degrees or the PIP joint,  
11 20 degrees.

12           This is done in my practice under a regional  
13 anesthetic, but can also be done under a local  
14 anesthetic with epinephrine. It's done through a  
15 extensile approach. I typically prefer an excision  
16 which kind of zigzags up the palm so that we can fully  
17 dissect out the diseased tissue. As we mentioned  
18 earlier, especially as we get up into the finger, the  
19 diseased cord which is seen here can be above the  
20 neurovascular bundle which is seen along right here,  
21 and kind of spiral around it.

22           So we need to meticulously dissect out the

1 nerve and artery, separate that from the diseased  
2 cord, and then ultimately, we'll excise that cord  
3 where it attaches to the flexor tendon sheath.

4           After the cord is removed, we then test our  
5 results. You know, oftentimes with the MP joint,  
6 we're able to get a full extension after excision of  
7 the diseased tissue. The PIP joint, however, doesn't  
8 always behave as well, and when there is a  
9 long-standing and high severe contracture, oftentimes  
10 we may still have a limitation of extension at the PIP  
11 joint level.

12           It's a matter of debate in the hand surgery  
13 literature of whether it's then beneficial to go in  
14 and formally release the ligaments about the PIP joint  
15 in order to obtain a better correction, or to try to  
16 achieve the rest of the correction postoperatively  
17 through therapy.

18           We get patients into therapy very quickly  
19 postoperatively because we don't want them to lose the  
20 ability to close their fist, which wasn't a problem to  
21 start with. Typically after surgery, particularly  
22 with patients who had a very severe contracture, we

1 may not be able to close all their skin incisions, and  
2 areas in the palm may be left open to heal. So if  
3 therapists will get involved earlier on in order to  
4 help manage the swelling that we always see after  
5 surgery, we want to minimize that, because that will  
6 limit the patient's ability to move their fingers, we  
7 want to start wound care if necessary; and we want to  
8 start those range of motion exercises.

9           Typically, I have patients in a splint  
10 full-time after surgery for the first two to four  
11 weeks. We kind of leave them in that splint to  
12 maintain that extended posture, and have them take it  
13 out of the splint every hour or two to work on their  
14 exercises. Once they can comfortably make a fist  
15 during the daytime, I have them just wear their splint  
16 at nighttime for approximately four months so that the  
17 scar that's formed after surgery doesn't contract at  
18 all and you don't see a recurrent contracture, or  
19 limit the recurrent contracture that we see.

20           This is just an example of a patient who's  
21 two days postop, who's moving her fist. This is the  
22 same patient who on the previous slide we had fully

1 extended her fingers, and you can see that she's not  
2 able to do that actively. Oftentimes, with advanced  
3 contracture of the PIP joint, the extensor tendons may  
4 be a little bit loose, they may be a little kind of  
5 bound down, may not have the strength to fully open on  
6 their own, which is why that therapy's so important.

7           This is a typical series of subtotal palmar  
8 fasciectomy. This is a consecutive series of 109  
9 patients in 2007. And what they found is that with  
10 the MP joint, they had a 97, 98 percent initial  
11 result. At the PIP joint, it was in the 70 percent  
12 range. And when they stratified it by severity, they  
13 found that the patients with a low severity  
14 contracture, less than 30 degrees, 78 percent  
15 maintained their correction at a year. However,  
16 patients with a more severe contracture which was  
17 greater than 60 degrees, only 50 percent of them  
18 maintained their correction that was achieved  
19 interoperatively.

20           Complications with surgery include digital  
21 nerve and artery injuries, particularly in recurrent  
22 cases; flare reaction, which is similar to complex

1 regional pain syndrome where the whole hand will  
2 become swollen and stiff; infection, loss of the  
3 ability to make a full fist and recurrence. And in  
4 this study, there was an average of about 20 percent  
5 of patients who had recurrent disease at 12 months.

6           So surgery has some limitations. The  
7 incision and dissection that's required to do the  
8 procedure safely leads to postoperative pain, healing  
9 response and scar tissue formation. Patients  
10 typically require a minimum of six weeks for their  
11 scars to settle down, and oftentimes three to four  
12 months. Hand therapy has been showed to optimize  
13 results. There are complications. It doesn't cure  
14 the disease and recurrence can still occur, and it's  
15 an operation that not every patient is willing to  
16 endure.

17           I find it helpful when talking to patients  
18 with Dupuytren's disease or any hand condition,  
19 discuss the options with them and to keep these goals  
20 in mind. We want to eliminate their contracture. We  
21 want to maintain a supple finger for the patients so  
22 they can comfortably open and close their fist. We

1 want to limit the morbidity that they go through,  
2 limit recurrence, limit complications and get them  
3 back to function as quickly as possible.

4 I'm excited to be here today as the  
5 Committee considers a new, novel option for  
6 Dupuytren's disease which will hopefully give us more  
7 options for our patients.

8 I'd like to now bring up Dr. DelConte.

9 DR. DELCONTE: Thank you, Dr. Kaplan.

10 My name is Tony DelConte, and I'm Auxilium's  
11 chief medical officer. And what I'd like to do this  
12 morning is discuss the overall clinical program and  
13 clinical efficacy for AA4500.

14 The clinical development program consisted  
15 of 13 studies in over 1,000 subjects who received at  
16 least one injection of the .58 milligram dose. These  
17 were done in a series of standard Phase 1, Phase 2,  
18 which included proof of concept and dose ranging, and  
19 then Phase 3 studies which we included as  
20 investigators orthopedic hand surgeons, plastic  
21 surgeons and rheumatologists.

22 In the Phase 3 study, there were three

1 double-blind placebo-controlled studies, and these  
2 were all followed by an open-label extension. And we  
3 had additional open-label studies and supportive  
4 studies for our safety database.

5           Now, if we turn first to the PK results.  
6 This is a series of 16 subjects with Dupuytren's  
7 disease who each received one injection, a single  
8 injection of .58 milligrams. And sampling was done at  
9 baseline and then at least 11 different time points  
10 through a 30-day period, and at no time point was any  
11 quantifiable systemic exposure noted, indicating that  
12 this is local, nonsystemic therapy.

13           Since the three double-blind  
14 placebo-controlled trials were all identically  
15 designed, I'll describe them here. A dose of .58  
16 milligrams or placebo was injected into the cord, into  
17 the pathologic structure, at each injection cycle.  
18 And a cycle consisted of the injection at day zero,  
19 and this was followed by the finger extension or  
20 manipulation procedure to disrupt the cord 24 hours  
21 following the injection.

22           And then further evaluations were done, and



1 then finally, at Day 30, an evaluation was done and  
2 measurements were done to see if the patient would be  
3 eligible to receive an additional injection. And each  
4 patient in the trial can receive up to three  
5 injections at four-week intervals, and this is the  
6 goal to achieve the primary outcome, the primary  
7 endpoint is a reduction in contracture to zero to 5  
8 degrees. That's to get the hand perfectly extended.

9           And each of the double-blind components of  
10 the trials were then followed by an open-label  
11 extension to allow patients on placebo to receive  
12 active drug.

13           The key inclusion criteria, these were  
14 adults at least 18 years of age who were affected with  
15 Dupuytren's disease and a palpable cord, causing a  
16 contracture of at least 20 degrees. And for the MP  
17 joints, they can go up to 100 degrees. For PIPs, this  
18 would be up to 80 degrees.

19           We excluded patients with bleeding disorders  
20 or disorders affecting the hand or any other condition  
21 that could confound the results. They could not have  
22 received previous treatment within three months prior

1 to the study start, and we excluded a few certain  
2 drugs and allergies to collagenase or any of the  
3 components of the product.

4           The efficacy assessments were done as  
5 follows: We measured the hand and the fingers at full  
6 extension and then full flexion, and the difference  
7 between flexion and extension was then recorded as the  
8 range of motion. We used an instrument like this,  
9 which is known as a goniometer, and this would be  
10 complete extension, and then a contracture of 90  
11 degrees would be to here. And these were done  
12 consistently on all of the subjects in the trial.

13           The patients were then randomized two to  
14 one, active to placebo, and there was further  
15 stratification done by the joint type, whether these  
16 were MP or PIP, and in Studies I and II, also by  
17 baseline severity. So we looked at low versus high  
18 severity.

19           Standard safety assessments were done,  
20 including the recording of adverse events, antibodies,  
21 standard laboratory and vital signs.

22           Now, the primary endpoint, the primary

1 outcome of all of the studies, was the proportion of  
2 subjects who achieved that correction to within zero  
3 to 5 degrees after their last injection, and this was  
4 defined as "clinical success" in the protocol. And  
5 there were multiple supportive secondary endpoints  
6 that were evaluated as well, and this was the  
7 proportion of subjects who achieved at least a 50  
8 percent reduction in their contracture angle. We  
9 considered this "clinical improvement." And then the  
10 percent change from baseline of the contraction angle  
11 was measured. We also evaluated time to success and  
12 the change in range of motion.

13           Additionally, there were global assessments,  
14 both physician and patient assessments done, to get an  
15 overall picture of the success of the therapy.

16           And here are the demographics and the  
17 disposition of the subjects. In the three double-  
18 blind placebo-controlled studies, A57, A59 and 303,  
19 which we refer to as Studies I, II and III, more than  
20 90 percent of the patients completed all of the  
21 assessments that were required by the protocol. And  
22 there was a predominance of men over women in the

1 studies, and the average age was about 62 to 63. And  
2 this is common of what you might see in a population  
3 of Dupuytren's patients who present for treatment.

4           And here are the primary endpoint results in  
5 all of the three studies. On the vertical axis is the  
6 proportion of patients who achieved success. That's  
7 the zero to 5 degrees. And what you see is in the  
8 three double-blind placebo-controlled trials, all of  
9 them met the primary endpoint and had a greater number  
10 of patients on drug versus placebo, where you had very  
11 few of the patients. In this largest study, 64  
12 percent on active versus just about 7 percent on  
13 placebo.

14           There were a series of secondary endpoints  
15 that were done in a hierarchical fashion, and I'd like  
16 to take you through a roadmap of how the secondary  
17 endpoints were done. Now, the primary endpoint was  
18 the reduction in contracture. But each of these was  
19 then taken for all joints first, and we looked at  
20 clinical improvement, then 50 percent reduction,  
21 percent change, time to reduction, change in range of  
22 motion. And these made up Secondary Endpoints 1

1 through 4.

2           This sequence was then repeated for the MP  
3 joints, and this made up Secondary Endpoints 5 through  
4 9. Again, repeated for PIP, 10 through 14. And this  
5 whole series was repeated again not after the last  
6 injection, but just after a single injection, and that  
7 made up Outcomes No. 15 through 26.

8           And if we then look at all of this together,  
9 we see the three studies and all of the secondary  
10 endpoints listed here. And in Study II, we're able to  
11 achieve nine additional of these secondary endpoints;  
12 Study III, most of the endpoints that were measured  
13 were achieved, but Study 1 hit the primary endpoint in  
14 all 26 of the secondary endpoints as I had described.

15           We also looked at the angle or degree of  
16 contracture, both before and after therapy for each of  
17 the three studies. In Study I, the patients started  
18 off about 50 degrees before therapy. And, again,  
19 referring to the goniometer, a 50-degree contracture  
20 would be to about here. Following therapy, the  
21 average contracture was about 12 degrees, or about  
22 here.

1           And then in placebo, they started off about  
2   the same place, around 49 degrees, but there was  
3   minimal effect on placebo changing contracture. And  
4   we see similar results for Study II as well as Study  
5   III in terms of fixed flexion contracture.

6           If we look at range of motion, an important  
7   functional parameter, we see that in Study I and Study  
8   II, patients started off with a range of motion going  
9   through an arc of a little over 40 to 45 degrees. But  
10   after therapy, this increased by almost 37 degrees and  
11   35 degrees in Study II, which was statistically  
12   significant over placebo. Minimal changed noted in  
13   the placebo group.

14           Now, I mentioned we evaluated the patient  
15   and physical global assessments. We looked at  
16   treatment satisfaction on a five-point analogue scale.  
17   And in this slide for Study I, the percentage of  
18   patients who had these results, 87 percent were either  
19   very or quite satisfied. And this was statistically  
20   significant from placebo, where most of the patients  
21   were in the very dissatisfied group. We did the same  
22   thing for a physician global assessment of the overall

1 treatment. And in this seven-point analogue scale, 85  
2 percent of the physicians rated the active treatment  
3 as either very much or much improved, compared to 93  
4 percent in the placebo that had no change.

5           We looked at the durability and recurrence  
6 rates. In all of the studies combined, there were 830  
7 successfully treated joints that met the primary  
8 endpoint. Thirty of these, or about 4 percent, had a  
9 recurrence of contracture, and this is after follow-up  
10 in some of the patients beyond one year. Half of  
11 these occurred between about three to six months of  
12 follow-up, and the mean follow-up period was a little  
13 over seven months.

14           To further assess the long-term follow-up of  
15 the recurrence and the durability, we are conducting a  
16 follow-up study in all of the patients who were  
17 enrolled in the trials who had improvement, and then  
18 to see what happens to their contractures after long-  
19 term. We also will be assessing the progression of  
20 disease in patients who either did not receive  
21 treatment or did not have success or a measurable  
22 improvement.

1           So to summarize the efficacy, all of the  
2 double-blind studies met the primary endpoint, and  
3 that is, more patients on AA4500 achieved this  
4 reduction to zero to 5 degrees over placebo. There  
5 were multiple supportive secondary endpoints,  
6 including improvement in range of motion, which  
7 support the efficacy. And both physician and patient  
8 satisfaction was significantly better for the drug  
9 over placebo. And overall, this provides efficacy  
10 comparable to what we see with surgical correction.

11           I'd now like to bring up Dr. Jim Tursi, who  
12 will discuss the clinical safety and the risk  
13 management activities that are proposed.

14           DR. TURSI: Thank you, Tony.

15           My name is Jim Tursi, and I'm the vice  
16 president of Clinical Affairs for Auxilium  
17 Pharmaceuticals. You've had an opportunity to witness  
18 the demonstrated efficacy profile. Now, we'd like to  
19 provide you a comprehensive view of the safety  
20 profile, and then I'll follow that by a very detailed  
21 look at our proposed risk management activities.

22           So first considering the safety profile,



1 I'll begin with an overview of our safety database.  
2 We'll consider subject disposition, extent of exposure  
3 as well as duration of follow-up. Then I'll speak to  
4 the adverse event profile. We'll consider local  
5 adverse events, serious adverse events, as well as  
6 those additional safety parameters. And lastly, as a  
7 biological, I'll speak to the immunologic response to  
8 AA4500.

9           Our pooled safety population is made up of  
10 1,082 subjects that were drawn across studies in our  
11 clinical program. They ranged from Phase 1 through  
12 Phase 3 and included both double-blind  
13 placebo-controlled trials as well as open-labeled  
14 studies.

15           The disposition of the 1,082 subjects  
16 includes a completion rate of 87.6 percent. 12.4  
17 percent discontinued, with the most common reasons:  
18 lost to follow-up and withdrawal of consent. Now, the  
19 subject age range was quite broad, and it ranged from  
20 age 33 to age 90. And subjects may have received  
21 anywhere from one to up to eight injections.

22           In terms of the extent of exposure, that

1 1,082 subjects represents 2,630 injections. That  
2 reflects treatment of 1,780 cords, and that's divided  
3 into 1,036 metacarpophalangeal cords and 743 proximal  
4 interphalangeal cords.

5 As to duration of follow-up, the mean  
6 duration, 9.5 months, with a minimum of two days and a  
7 maximum of 6.7 years. The interjection interval, time  
8 between injections, ranged from as short as ten days  
9 to as long as greater than 6.4 years.

10 Next, to the adverse event profile, and as  
11 we discuss this, I would ask you to consider the acute  
12 and nonsystemic nature of AA4500 therapy. When  
13 considering those adverse events that occurred at  
14 greater than or equal to 5 percent, the most common:  
15 edema peripheral or swelling of the treated hand,  
16 contusion and injection site pain, were the three most  
17 common. And they ranged from 77 percent to 40.9  
18 percent. The vast majority of these were mild to  
19 moderate in severity, with less than 3 percent being  
20 considered severe.

21 The next most common adverse events:  
22 extremity pain, injection site hemorrhage, tenderness,

1 injection site swelling, ecchymosis and skin  
2 laceration. And that ranged from 37.4 percent to 12.7  
3 percent. Again, the vast majority mild to moderate in  
4 intensity, with less than 1 percent of these adverse  
5 events being considered severe.

6 Finally, completing those to greater than or  
7 equal to 5 percent pruritus, lymphadenopathy, blood  
8 blister, axillary pain, hematoma, arthralgia and  
9 injection site pruritus, ranging from 12.7 percent to  
10 5.3 percent. Again, the vast majority were mild to  
11 moderate, with less than one-half of 1 percent of this  
12 adverse events being considered severe.

13 There are several important trends to bring  
14 forward as it relates to the adverse event profile.  
15 The overwhelming majority of adverse events were  
16 confined to the treated extremity. Most were  
17 nonserious and were either of a mild or moderate  
18 intensity. The vast majority resolved prior to the  
19 next injection with no further intervention, with a  
20 median duration across the entire adverse event  
21 profile of ten days.

22 Next, considering serious adverse events in

1 the clinical program. There were 92 subjects who  
2 experienced serious adverse events. But it's  
3 important to point out that if the serious adverse  
4 event did not involve the treated extremity, there was  
5 a similar proportion between AA4500 subjects and  
6 placebo subjects. Nine subjects experienced ten  
7 serious adverse events that were considered treatment-  
8 related. They included a case of ligament injury,  
9 three cases of flexor tendon rupture, a recurrent case  
10 of complex regional pain syndrome, a boutonniere  
11 deformity, a case of deep vein thrombosis of the lower  
12 extremity, a case of sensory disturbance and  
13 Dupuytren's contracture in the same subject, and a  
14 case of tendonitis.

15 I would like to spend some time and speak  
16 specifically and provide details around the case of  
17 ligament injury and the flexor tendon ruptures. The  
18 first case was a 61-year-old male who, 43 days  
19 following his second injection, noted on his physical  
20 exam significant bow stringing. Essentially, the  
21 flexor tendons were pulling forward on the skin of the  
22 treated finger. He was ultimately diagnosed with an

1 A-2 and an A-4 pulley rupture, and surgical correction  
2 in the form of joint fusion and tenotomy was  
3 ultimately performed.

4           The second case, a 62-year-old male who, six  
5 days following his first injection, noted finger  
6 weakness. Physical exam and MRI confirmed a rupture  
7 of the flexor digitorum superficialis tendon, with an  
8 intact flexor digitorum profundus tendon. This  
9 subject had a pre-existing boutonniere deformity, and  
10 it was ultimately brought to surgical correction of  
11 that deformity with no surgical intervention of the  
12 tendon rupture at that time.

13           The third case was a 61-year-old male who,  
14 eight days following his first injection, had resumed  
15 full normal activities. This included his employment,  
16 which required him to lift heavy objects. During  
17 employment while lifting a heavy pallet, he noted  
18 immediate finger swelling and weakness, and  
19 ultimately, MRI and physical exam confirmed a rupture  
20 of the flexor digitorum profundus tendon and a partial  
21 tear of the flexor digitorum superficialis. The  
22 subject underwent tenolysis as repair.

1           The fourth case was a 76-year-old male who,  
2   four days following his third injection, noted an  
3   inability to flex the treated finger. Physical exam  
4   confirmed rupture of both the FDS and FDP tendons, and  
5   ultimately, a two-stage repair with tendon grafting  
6   procedure was performed.

7           Although these four events represent less  
8   than one-half of 1 percent of the safety population,  
9   it's clearly important to understand the anatomy of  
10   Dupuytren's disease and the underlying flexor tendons.  
11   This photograph was taken from an operative correction  
12   of a Dupuytren's cord. And just to point out the  
13   anatomy, the Dupuytren's cord in white, and at the  
14   base of the ruler is the intact flexor tendon. But  
15   the point being that there are areas where the cord is  
16   in close proximity to the tendon, and other areas  
17   where it is more distant.

18           These four cases were considered to the  
19   effect of AA4500, and as such, is a focus of the risk  
20   management plan which I'm going to discuss in just a  
21   few moments.

22           What about additional safety parameters? We

1 checked laboratory values, including renal function  
2 and liver function studies, and the percentage of  
3 subjects in the AA4500 group with abnormalities was  
4 low, and was comparable to that in the placebo group.  
5 In terms of hematology parameters, again, the percent  
6 of subjects in the AA4500 group was low with  
7 abnormalities, and was comparable to the placebo  
8 group.

9           And finally, we also checked vital sign  
10 parameter changes, blood pressure, heart rate,  
11 respiratory rate. The number of subjects with  
12 clinically meaningful changes was low, and was  
13 comparable to the placebo group.

14           As a biologic product, we would expect to  
15 see potentially an immune or an immunologic reaction  
16 to treatment with AA4500. First, considering subjects  
17 who received a single dose, and to orient you: across  
18 the vertical axis is the mean log titer of antibodies,  
19 of either anti-AUX-I or anti-AUX-II in green and  
20 orange respectively; and across the horizontal axis,  
21 the time in months after injection.

22           And what was demonstrated for both anti-AUX-

1 I and anti-AUX-II was a peak in antibody titer at  
2 approximately two to four months, with a waning  
3 thereafter.

4           When considering subjects who received  
5 multiple injections, in this case up to eight  
6 injections, again on the vertical axis, the mean log  
7 titer, across the horizontal axis, the respective  
8 injection number. First, when considering anti-AUX-I,  
9 we see an increase in antibody titers that essentially  
10 peaks at about the fifth or sixth injection, and then  
11 plateaus thereafter. For anti-AUX-II, again, we see a  
12 similar pattern, an increase in antibody titer through  
13 about the fifth or sixth injection, with a plateau  
14 thereafter.

15           As to seropositivity, the percentage of  
16 subjects who have antibodies present, by the third or  
17 fourth dose, 100 percent of subjects have antibodies  
18 present to anti-AUX-II or anti-AUX-I respectively.

19           Considering that virtually 100 percent of  
20 subjects develop antibodies, the question becomes do  
21 these antibodies affect the safety profile of AA4500.  
22 So we performed multiple analyses, including examining



1 the rate, the severity and the duration of the adverse  
2 event profile. In addition, we looked for evidence of  
3 systemic anaphylactic reactions.

4           So first, to consider the rate, if anti-drug  
5 antibodies were to negatively affect the safety  
6 profile of AA4500, we would expect the rate of adverse  
7 events to consistently increase with increasing  
8 antibody titers. When considering the four most  
9 common adverse events, across the vertical axis is the  
10 percentage rate of the specific adverse event, which  
11 is identified above each table, and across the  
12 horizontal axis by injection number. And what's  
13 demonstrated is with increasing antibody titers and  
14 increasing injections, there is no consistent pattern  
15 of increasing adverse events rates with subsequent  
16 injections, and thus with increasing antibody titers.

17           This profile was consistent across the  
18 entire adverse profile of AA4500, and demonstrates  
19 that there was no consistent pattern between adverse  
20 rates and increasing antibody titers.

21           Then we considered severity of the adverse  
22 events. If anti-drug antibodies were to negatively

1    affect the safety profile, we would expect those  
2    subjects with severe adverse events to have higher  
3    antibody titers. To orient you: across the vertical  
4    axis is the mean line titer of either anti-AUX-I in  
5    green or anti-AUX-II in orange, and across the  
6    horizontal axis are the cohorts of those that did not  
7    experience the adverse event -- in this case, it's  
8    swelling of the hand -- or experience the adverse  
9    event as mild, moderate or severe.

10            So when we first consider those subjects who  
11    did not experience swelling of the hand, the mean log  
12    titer was 3.5. When we then look at subjects who  
13    experience the adverse event as mild, moderate or  
14    severe, it's clear there's no correlation between  
15    adverse event absence or presence, or no correlation  
16    between the severity of the adverse event and the  
17    antibody titer. That was also found for anti-AUX-II.

18            Considering those four most common adverse  
19    events, contusion, no correlation; injection site  
20    pain, again, no correlation; and extremity pain with  
21    no correlation. This lack of correlation was  
22    demonstrated across the entire adverse event profile,

1 confirming that adverse event severity does not  
2 correlate with antibody titer.

3           Then we looked at the duration of adverse  
4 events. Should anti-drug antibodies negatively affect  
5 the safety profile, we would expect the duration of  
6 adverse events to increase with increasing antibody  
7 titers and subsequent injections. Across the vertical  
8 axis, the median duration is days; across the  
9 horizontal axis, the injection number. And, again,  
10 as is demonstrated, there is no consistent increase of  
11 adverse event duration with subsequent injections.  
12 And, again, these findings were across the entire  
13 adverse event profile for AA4500.

14           Now, that confirmed that the duration of  
15 adverse events does not correlate with subsequent  
16 injections and increasing antibody titers.

17           Next, we did a thorough evaluation to look  
18 for any signs, symptoms or signals of systemic  
19 anaphylaxis reactions in the clinical program. And  
20 that was very straightforward. There were none in the  
21 clinical program.

22           So in summary, with a safety database of

1 nearly 1,100 subjects and an injection database  
2 representing over 2,600 injections, the most frequent  
3 adverse events were confined to the treated extremity.  
4 They were either mild or moderate in intensity, with  
5 the vast majority resolving prior to the next  
6 injection.

7           Serious adverse events occurred, including  
8 tendon rupture and ligament injury, and that risk is  
9 clearly identified and will be a focus of our risk  
10 management plan, which I'm going to discuss in much  
11 more detail in just a few minutes.

12           As it relates to routine laboratories and  
13 vital signs, there were no clinically meaningful  
14 differences demonstrated between AA4500 subjects and  
15 placebo subjects. As to immunogenicity, antibodies  
16 developed in nearly all subjects, but they do not  
17 appear to adversely affect the safety profile.

18           And finally and importantly, there were no  
19 events or signals indicative of systemic anaphylaxis  
20 in the clinical program.

21           In order to ensure that our clinical trial  
22 results are accomplished in clinical practice, we've

1 created a risk management plan which we believe is  
2 comprehensive and will be effective in that endeavor.  
3 In order to do that, first we must lay out several  
4 goals of that risk management plan: first and  
5 foremost, to ensure appropriate administration of  
6 AA4500. In order to do that, we must recognize  
7 potential and identified risks. We must create and  
8 implement strategies ultimately to minimize those  
9 risks, and we must inform and educate both physicians  
10 and patients.

11 First considering the potential and  
12 identified concerns. Clearly, injected-related  
13 bleeding in subjects with coagulation disorders would  
14 be a risk of any injection therapy, and thus is a  
15 potential risk of AA4500 treatment. The potential for  
16 allergic reaction with a biological is also a  
17 potential risk. Identified tolerability and safety  
18 concerns include those localized reactions which I've  
19 provided some detail around as well, as the risk of  
20 tendon rupture and ligament damage.

21 As to those potential risks, risk management  
22 activities would primarily include labeling of the

1 product to address these concerns. Injection-related  
2 bleeding in subjects with coagulation disorders would  
3 be an expected risk of an injectable therapy, so the  
4 label will include a caution for use in those with  
5 coagulation disorders. Use would not be recommended  
6 for those on concurrent anticoagulant medications, and  
7 consistent with the clinical program, however,  
8 prophylactic low dose aspirin use would be considered  
9 acceptable.

10           As to the potential risk of allergic  
11 reaction, the label would include a contraindication  
12 for use in any individual with a known  
13 hypersensitivity to AA4500. And consistent with most  
14 medications, it would include a warning to physicians  
15 to prepare to address any potential allergic reactions  
16 should they occur.

17           As to the identified tolerability and safety  
18 concerns, I first spoke of the localized reactions.  
19 They are common and they're expected with AA4500  
20 treatment. You've heard the most common: edema  
21 peripheral, swelling of the treatment hand, bruising  
22 and injection site pain. While the vast majority were

1 mild to moderate with resolution prior to the next  
2 injection, clearly, it's essential that both  
3 physicians and their patients know what to expect with  
4 treatment from AA45.

5 Risk management activities as to the local  
6 reactions will include product labeling, physician  
7 training and patient product information. The product  
8 labeling will clearly describe the local reactions.  
9 Consistent with the clinical programs, multiple cords  
10 should not be treated simultaneously, and only one  
11 hand should be treated per session.

12 Physician training, which I'll go into much  
13 more detail in just a few minutes, will include  
14 details of these local reactions. So physicians  
15 during the training period prior to use of AA4500 can  
16 know what to expect regarding these local reactions.

17 And patient product information will  
18 describe these local reactions in easy-to-understand  
19 and detailed language so patients can know what to  
20 expect before, during and following therapy with  
21 AA4500.

22 While the four cases of tendon rupture

1 and/or ligament rupture represented less than one-half  
2 of 1 percent, clearly, inappropriate exposure to  
3 normal collagen-containing structures can result in  
4 lysis of collagen and subsequent to damage to those  
5 structures ultimately resulting in possible injury or  
6 reduction of functionality.

7           The risk management plan is quite  
8 comprehensive as it pertains to this specific risk.  
9 It will include product labeling aspects -- and I'll  
10 go into each of these in quite some detail --  
11 physician training and access management program,  
12 safety monitoring which is enhanced; and patient  
13 education.

14           So first focusing on product labeling, the  
15 product labeling will be quite detailed and very  
16 informative for physicians. The intended users of  
17 AA4500 are physicians experienced in the diagnosis and  
18 management of Dupuytren's disease: hand surgeons,  
19 orthopedic surgeons, plastic surgeons, general  
20 surgeons with a hand focus and rheumatologists.

21           The risk of tendon rupture will be clearly  
22 identified, and an injection precaution is also



1 included. And that reads, "Because AA4500 lyses  
2 collagen, care should be taken to avoid injecting into  
3 normal collagen-containing structures of the hand.  
4 Exposure of collagen-containing structures to AA4500  
5 may result in damage to their structures and possible  
6 permanent injuries such as tendon rupture or ligament  
7 damage."

8           As you can see, it's quite detailed, and  
9 physicians experienced in this disease would clearly  
10 understand the warning as it is written.

11           The next component will be physician  
12 training, and we believe that physician training is  
13 essential for a successful transition from clinical  
14 development to clinical practice. It's first  
15 worthwhile to consider the challenges that we face in  
16 our clinical program, to provide a little bit of  
17 history as to the clinical development program.

18           This was a new therapeutic procedure for  
19 Dupuytren's disease. There was very limited  
20 experience with AA4500 in this indication, and we were  
21 embarking on a multinational Phase 3 program. So we  
22 needed to essentially create a training program which

1    could be extrapolated from the experience of a very  
2    small number of physicians, and ultimately be able to  
3    extrapolate that to multiple investigators and  
4    multiple sites across multiple countries.

5                So we provided several injection training  
6    options for investigators. The first option was a  
7    30-minute injection training workshop in which some  
8    PowerPoint slides were reviewed, as well as a section  
9    of our injection training DVD for investigators. The  
10   second option was approximately 30 minutes of  
11   injection training at the investigator meeting, again  
12   composed of PowerPoint slides and a section of our  
13   injection training DVD. All clinical trial sites and  
14   investigators received a copy of our injection  
15   training DVD as well as our injection training manual.

16               What we found was there was some variability  
17   as to the preferred method of training for both  
18   primary and sub-investigators. So when we consider  
19   first the primary investigators -- and this focuses on  
20   Studies 857 and 859 -- of the 21 primary  
21   investigators, five attended both the injection  
22   training workshop and the injection training portion

1 of the investigator meeting. Four attended just the  
2 injection training workshop, and five attended just  
3 the injection training portion of the investigator  
4 meeting.

5           What was evident was the majority of primary  
6 investigators attended neither the injection training  
7 workshop or injection training at the investigator  
8 meeting. All had access to the injection training DVD  
9 or manual, with one primary investigator having an  
10 opportunity to directly observe a procedure.

11           As it relates to the sub-investigators, none  
12 attended the injection training workshop, two attended  
13 injection training at the investigator meeting, and,  
14 again, all had access to the injection training DVD or  
15 manual, with some having an opportunity to observe  
16 from the primary investigator.

17           It was clear when we spoke with them that  
18 they preferred utilizing the injection training DVD  
19 and the injection training manual. We confirmed that  
20 by meeting with not only our investigators but other  
21 practicing physicians, and these included hand  
22 surgeons, orthopedic surgeons, plastic surgeons and

1     rheumatologists.

2                 We reviewed what we had done previously in  
3     training methodology, and we specifically asked their  
4     advice, discussing their needs and their preferences.  
5     And overwhelmingly, they requested a video (and)  
6     written training program. They asked that it be clear  
7     and comprehensive, informative and accessible, and  
8     expanded from the clinical program.

9                 So in order to do that, we created a  
10    training program that is broader in scope and content  
11    than that which we used to train our investigators.  
12    The proposed program will include additional  
13    information to help physicians use AA4500  
14    appropriately. It will provide more depth, more  
15    examples, more animations and demonstrations based on  
16    the experience of our clinical investigators, and  
17    completion of training with attestation will be  
18    mandatory prior to accessing AA4500.

19                The training program is composed of an  
20    injection training DVD and injection training manual  
21    and the program components, anatomy and pathology,  
22    product preparation, injection, finger extension, a

1 frequently asked questions section, and self-  
2 assessment questionnaires. This was created with and  
3 features demonstrations of appropriate use by  
4 physicians with experience using AA4500. A hard copy  
5 training manual is also available for those that  
6 prefer that method of interaction or training.

7           In terms of the first component, review of  
8 anatomy and Dupuytren's pathology, this will include  
9 detailed illustrations to help the physician visualize  
10 the relationship between the Dupuytren's cord and  
11 other normal hand structures. It will include  
12 information on disease progression, as well as  
13 information regarding the mechanism of action of  
14 AA4500, so physicians can better understand the  
15 treatment procedure.

16           The demonstration of injection and the  
17 finger extension demonstrations include details on  
18 product preparation, needle placement advice specific  
19 to the joint being treated, details around the  
20 injection procedure, as well as a detailed description  
21 of the extension procedure, with a visualization of  
22 cord rupture.

1           The frequently asked questions section  
2 includes questions that are both product- and  
3 procedure-specific; questions regarding preparation,  
4 injection and finger extension.

5           In addition, potential and identified risks  
6 are discussed as part of the training program,  
7 including those local reactions we talked about and  
8 the identified risks of tendon rupture.

9           Also, information will be provided to  
10 physicians to ease adverse event reporting,  
11 essentially instructions to physicians during training  
12 as to how to report adverse events to Auxilium.

13           Lastly, a self-assessment questionnaire will  
14 be included to ensure physician understanding of  
15 content.

16           I would like to show you some excerpts.  
17 First, an excerpt from the injection technique  
18 section, and what I would ask you to do, realize this  
19 is a small excerpt of the draft version of the  
20 training materials, and it's intended for clinical  
21 practice, so you will hear a reference to the word  
22 "Xiaflex," our proposed trade name. In addition, I

1 would ask you to look for the detail, clarity,  
2 animation and live representation that's in the video.

3 (Video played.)

4 DR. TURSI: As you can see, it's quite  
5 detailed. It includes animation and live  
6 representation. Now I would like to show an excerpt  
7 from our extension procedure video. This includes  
8 information regarding those local reactions, as well  
9 as details of what physicians can expect during the  
10 extension procedure.

11 (Video played.)

12 DR. TURSI: As was demonstrated in the  
13 video, with complete correction of the hand in this  
14 patient was that audible pop, providing physicians  
15 with knowledge as to what to expect. What's also  
16 evident is some context around these local reactions:  
17 bruising, swelling of the hand and contusion.

18 We believe training will be most effective  
19 if it's required in order to access AA4500, and that's  
20 the intent of the access management program. Training  
21 will be required to access AA4500 by physicians  
22 experienced in the diagnosis and management of

1 Dupuytren's disease. They must attest to completion of  
2 the injection training video or manual, and  
3 ultimately, they must submit attestation to Auxilium  
4 for enrollment in order to receive access.

5           Diagrammatically, if a physician wants to  
6 use AA4500 and they're not enrolled, they will be  
7 referred to physician training. That could be via  
8 website, directly via the video or training manual.  
9 With completion of training and attestation, they  
10 would forward their signed enrollment form to  
11 Auxilium, at which point, they would be placed in a  
12 central database of enrolled physicians. Once  
13 enrolled, they would contact their distributor  
14 requesting access to AA4500. The distributor would  
15 check the enrollment database to ensure that they're  
16 enrolled. If they're not enrolled, ultimately, they  
17 would be redirected for physician training, and  
18 ultimately, for enrollment. If they are enrolled,  
19 they would receive access to AA4500.

20           The next consideration would be an enhanced  
21 safety monitoring program, and that would be essential  
22 and vital to identify any potential safety signals.



1 Safety activities will include a safety hotline which  
2 will help ease case reporting for physicians. And as  
3 I noted, the training program will include information  
4 for physicians to improve and ease that reporting.

5 We'll perform an aggregate safety review by  
6 an Auxilium safety physician monthly for the first  
7 year, followed by quarterly reviews thereafter.

8 And in the event of a tendon rupture, we  
9 will follow up directly with the physician with a  
10 tendon rupture questionnaire. This is a draft  
11 version, and certainly, my intent is not to take you  
12 through each detail of the questionnaire, merely to  
13 provide you a view of the comprehensive nature of the  
14 questionnaire, the amount of information requested,  
15 and the details that are requested specifically of the  
16 document.

17 No risk management activities will be  
18 complete without considering the patients suffering  
19 from Dupuytren's disease. So we'll provide multiple  
20 portals for these patients to access information. It  
21 will include the patient product information leaflet,  
22 as I said, written in easy-to-understand language so

1 patients know what to expect before, during and after  
2 therapy. It will include web-based resources,  
3 information on the disease state, but also trained  
4 physician listings, so patients can determine in their  
5 region physicians who've attested to training with  
6 AA4500.

7           We'll provide office-based educational  
8 materials and a toll-free patient product information  
9 line for further questions.

10           I've described each of these individual  
11 pieces in some detail, but I think it's important to  
12 step back and consider the comprehensive nature of  
13 this plan, comprehensive to the needs of both  
14 physician and patient. It's constructed of many  
15 components, to build a strong foundation for the safe  
16 and effective use of AA4500 in clinical practice.

17           At the outset, I spoke of the goals,  
18 primarily to ensure appropriate administration of  
19 AA4500. We believe our risk management program is  
20 comprehensive and will be successful in this endeavor.  
21 To recognize those potential and identified risks, it  
22 creates and will implement strategies to minimize

1    those risks.  It educates and informs both physicians  
2    and their patients suffering from Dupuytren's disease.  
3    We believe it creates the optimum environment to  
4    transition AA4500 from clinical development to  
5    clinical practice.

6                    So thank you, and I would like to ask  
7    Dr. DelConte to come up for the final overall summary.

8                    DR. DELCONTE:  Thank you, Jim.

9                    Over the last hour, you've heard a lot about  
10   AA4500, and I do look forward to an active discussion  
11   with members of the Advisory Committee.  But before we  
12   go into that, first I would like to summarize why we  
13   believe that AA4500 should be approved as the first  
14   nonsurgical therapy for Dupuytren's disease.

15                   First, we've heard from Dr. Kaplan how  
16   Dupuytren's disease is a debilitating condition that  
17   affects everyday activities of those afflicted with  
18   the disease.  He explained to us that the first  
19   approach is often observation and reassurance.  Once  
20   the disease progresses to the point where the patient  
21   is willing to have surgery, the results are generally  
22   good.  Surgery can typically provide relief,

1     straighten joints and restore function.

2             However, surgery is not a perfect solution.  
3     While most surgeries have a positive result, there's  
4     some serious risks and complications that occur,  
5     including injury to other structures such as nerves  
6     and arteries. There's also risk of infection,  
7     scarring and general wound healing issues. In  
8     addition, the surgical procedures leave the patient  
9     with a prolonged follow-up and recovery period,  
10    sometimes requiring extensive physical therapy. And  
11    subsequent surgeries to the same area are more complex  
12    and involve more risk.

13            Turning to AA4500, we've demonstrated  
14    efficacy in three double-blind placebo-controlled  
15    trials, each of which met that stringent primary  
16    endpoint of getting to zero to 5 degrees, thus  
17    restoring function. And looking specifically at Study  
18    I data which was recently published this month in the  
19    New England Journal of Medicine, 64 percent of the  
20    patients achieved that primary endpoint compared to  
21    just under 7 percent with a placebo.

22            And secondly, the safety profile of AA4500

1 has been well-characterized, with most adverse events  
2 being local, self-limiting and confined to the treated  
3 extremity.

4           And thirdly, in order to generalize the  
5 results, we've developed a comprehensive training  
6 program that has been designed and modeled after our  
7 investigator training, and further enhanced to ensure  
8 that the clinical results seen in our trials can be  
9 extrapolated with an appropriate population of  
10 physicians and patients.

11           In summary, AA450 will provide the first  
12 nonsurgical therapy for managing Dupuytren's disease.

13           I thank the panel for your attention, and  
14 I'd like to join my colleagues now.

15           DR. O'NEIL: Thank you. We will now have a  
16 discussion of the data presented, with the panel  
17 asking questions of the sponsor. I would like to ask  
18 my colleagues on the panel to please signal a comment  
19 that they may have and wait for recognition by the  
20 Chair so we don't all talk together, and also remind  
21 you to turn off your microphone after you have spoken  
22 so we don't have sheer chaos and wild noise.

1                   I would like to begin with a question  
2   probably for Dr. Tursi, but also for the  
3   pharmacologist involved in the development of this  
4   product. Collagenase is one of a large family of  
5   enzymes in almost any living organisms that are in a  
6   class called serine proteases. These serine proteases  
7   are very potent and multifunctional enzymes that do  
8   more than what we have named them to do. In  
9   particular, the complement system is a series of  
10   serine proteases which work one upon another to  
11   activate enzymes that have large amplification and  
12   very broad complications when allowed to proceed  
13   uninhibited in the body.

14                  Collagenase, elastase, complement proteins,  
15   thrombin, the kinins are interrelated and one can  
16   activate another. There are a number of anti-  
17   proteases that control these reactions in the body.  
18   I'm wondering if in any of your animal development or  
19   in your human studies, you found evidence of  
20   complement activation, thrombin activation -- I  
21   noticed one of your SAEs was a DVT in a remote  
22   extremity -- or other related things.

1           I suspect a lot of the local edema is from  
2   kinin activation locally. Do we have any information  
3   about that, because this could -- if the product were  
4   injected near or worse still in a vessel, could  
5   certainly produce remote reactions.

6           DR. DELCONTE: Yes, I'd like to ask Dr.  
7   Susan Hart, who's our toxicologist, to come up, and  
8   she can describe some of the animal findings,  
9   including the histologic findings.

10          Dr. Hart.

11          DR. HART: I'll speak directly to your  
12   question regarding activation of complement or other  
13   serine protease pathways. We haven't evaluated these  
14   directly in animal studies because there is an  
15   extensive literature base on the effects of  
16   clostridial collagenases in these pathways. And  
17   having reviewed that literature, I found no evidence  
18   that the collagenases directly activate complement,  
19   directly convert kinin to bradykinin, or directly  
20   interfere with thrombin pathways or alter thrombin-  
21   mediated pathways.

22          We haven't seen any indication of that in

1 the animal studies which have included evaluation of  
2 coagulation parameters, hematology, local histology  
3 and also systemic histology. So as far as the  
4 literature is concerned and our own studies are  
5 concerned, there's no evidence that the product itself  
6 interferes with those pathways.

7 DR. TURSI: As to that specific adverse  
8 event, I can provide you a little bit more detail, but  
9 it did not appear to be related to AA4500 use. This  
10 was a 62-year-old male with a history of Lederhose  
11 consistent with a diathesis of Dupuytren's disease.  
12 And he was based in Australia and drove a considerable  
13 distance to the study site. This was approximately  
14 two to three hours in each direction. Had received  
15 the injection day zero and ultimately noted the lower  
16 extremity symptoms of left knee and calf pain two days  
17 thereafter. A Doppler revealed a single lower  
18 extremity thrombosis, and this was ultimately managed  
19 with anticoagulants.

20 Across the entire clinical program, there  
21 did not appear to be evidence of complement  
22 activation. In regards to the local events, it's



1   important to realize the pharmacology of AA4500 may  
2   also play a role ultimately in those local reactions.  
3   Anti-AUX-I and anti-AUX-II as enzymes are very  
4   efficient at cleaving collagen into small fragments.  
5   When they do so, especially in the animal studies, we  
6   see evidence of increased capillary permeability,  
7   hemorrhage, some rapid localized edema and local non-  
8   immunologic mass cell histamine release.

9               So a lot of the symptoms that we're seeing  
10   locally could also be explained by the pharmacology.  
11   And as I noted across the clinical database, there did  
12   not appear to be evidence consistent with your  
13   concern.

14              DR. O'NEIL:   Dr. Weisman.

15              DR. WEISMAN:   I have two questions.   I'm not  
16   sure which of you would address one or the other, but  
17   we'll just see.

18              The first question is, it seems that you've  
19   set up a very interesting, almost gatekeeper type of  
20   panel to authorize physicians to be able to use this  
21   procedure.   Who constitutes that panel?   How will that  
22   panel be independent of marketing efforts?   And that

1 panel would be somehow accountable to a review as to  
2 making sure that the review of these individuals who  
3 are allowed then or certified to be able to use the  
4 product continues on the very high level and is  
5 consistent with the collection of data about the  
6 results that you're also collecting of the procedure.  
7 How will that be arranged?

8 DR. DELCONTE: I'll have Dr. Tursi address  
9 the issue of the access management program, but the  
10 types of specialties was designed after the types of  
11 physicians who were in the clinical trial program.

12 DR. TURSI: Thanks. As I noted during the  
13 main presentation, the access management program's  
14 specific intent is to basically provide access to  
15 those physicians who are best-suited to ultimately use  
16 the product. One of the first steps ultimately in  
17 that access, as I noted, was the physician training  
18 component, and the required attestation of that  
19 training by the physician that would like to use the  
20 product.

21 To that end, attestation will require  
22 specifics that the physician identify their specialty.

1 If it's within one of those specialties we've  
2 described which was hand surgeon, orthopedic surgeon  
3 or plastic surgeon, rheumatologist, then the process  
4 would move quite automatically, through ultimately  
5 providing access to those physicians.

6           There also would be an opportunity for them  
7 to identify themselves as another specialty. If they  
8 do, that would then be called to the attention of our  
9 internal Auxilium staff, which likely would be through  
10 our safety group, at which time we would determine the  
11 availability for the drug for those individuals.

12           So the goal being to provide access  
13 ultimately to those physicians best-suited to use it,  
14 which would hopefully ultimately achieve the clinical  
15 trial results in clinical practice.

16           DR. WEISMAN: My second question is sort of  
17 a derivative of the first, and that is that since the  
18 complications that we're concerned about, such as  
19 tendon rupture or ligament rupture, and the fact that  
20 shortly after the procedure, many of these patients  
21 are going to require a manipulation for efficacy,  
22 there would need to be a great deal of expertise of

1 hand surgery involved either with the procedure itself  
2 or as a follow-up of the procedure.

3           So my question is a conceptual one, and that  
4 is, do you consider this a medical procedure or a  
5 surgical procedure, and should the individuals that  
6 are involved in this whole process be individuals who  
7 are specifically used to doing surgical-type  
8 manipulation and careful control of these factors  
9 rather than internists or rheumatologists who  
10 generally speaking are not used to doing these kind of  
11 procedures following an injection of this material?

12           DR. TURSI: We consider this a medical  
13 procedure, and we have in our group Dr. Kaplan, who's  
14 a hand surgeon who was an investigator, and Dr.  
15 Waller, who is a rheumatologist, and who also was an  
16 investigator in one of the open-label trials. So I'd  
17 like first, Dr. Kaplan, if you can discuss your view  
18 of the entire procedure, and then I'll have Dr. Waller  
19 come up as well.

20           DR. T. KAPLAN: I think as you mentioned,  
21 there are two main parts to the procedure, one putting  
22 the injection in place, and then secondarily, doing

1 the manipulation, which I agree is beneficial for  
2 probably most patients who don't rupture on their own  
3 spontaneously beforehand.

4 As far as doing an injection, it's fairly  
5 straightforward, I think, amongst both surgical  
6 specialties and rheumatology, internal medicine.  
7 Rheumatologists frequently do, to my knowledge, inject  
8 Dupuytren's cords. They do do cortisone injections  
9 for joints or trigger fingers as well in the hand. So  
10 I think that they're accustomed to doing injections  
11 even into Dupuytren's tissue. They may be less  
12 accustomed to doing manipulations, and I'll let Dr.  
13 Waller kind of address his experience with that.

14 I found the procedure relatively  
15 straightforward. As with any new procedure, there is  
16 some experience that you gain in the first couple  
17 times that you do, and certainly, I think that I've  
18 gotten better at it as I've done more of it. But,  
19 again, I think it is relatively straightforward, and I  
20 think it's something that would not be too difficult  
21 to teach or train to perform.

22 DR. WALLER: To reintroduce myself, Philip

1 Waller from Houston, Texas, practicing rheumatologist.  
2 I do think we've got the knowledge of the anatomy at  
3 least from tendons. We certainly have injected  
4 trigger fingers, Dupuytren's, de Quervain  
5 tenosynovitis, bicep tendonitis. This was obviously a  
6 different injection, and actually, almost a simpler  
7 injection in the sense that the cord was so different  
8 than what we've seen in joint and injecting other soft  
9 tissue.

10           The manipulation itself, I will agree it's  
11 not something we do every day in clinical practice.  
12 As Dr. Kaplan said, it was a learning process that  
13 after really with our first patient, it was a fairly  
14 simple procedure -- and certainly no disrespect to the  
15 hand surgeons or orthopedic surgeons, I do think we  
16 have the experience to do the manipulation, because it  
17 did not really require a specific amount of excess  
18 training. This video is a completely different video  
19 than we initially saw as an investigator, and much  
20 more comfortable to watch, in the sense that the  
21 training's much easier in this video.

22           So the answer, yes, I think we can do the

1 injection. Secondly, the manipulation I do believe  
2 can be done.

3 DR. O'NEIL: The next question is from  
4 Dr. Saag.

5 DR. SAAG: I want to follow up on Michael's  
6 comment about what types of providers should be  
7 performing this procedure, and tag on to the comment  
8 made by the hand surgeon that there's a bit of a  
9 learning curve. And particularly, as that relates to  
10 the risk management strategy, how can we be sure that  
11 by watching a video -- and for those in the room that  
12 have been asked to watch videos as part of training,  
13 unfortunately, many people are checking their e-mail  
14 at the same time while they're surfing on the web on  
15 the video. How do we assure that there is adequate  
16 knowledge and adequate experience gained just from  
17 this video, to avoid a significant learning curve?

18 And the corollary to that is, do we have any  
19 sense from the four cases of tendon rupture and  
20 ligament injury about where those events occurred in  
21 the experience of the investigator? And are we  
22 confident that the risk management program will

1 mitigate the potential for injecting this potentially  
2 toxic compound in areas where it's not supposed to be?

3 DR. DELCONTE: Dr. Tursi will answer that  
4 question.

5 DR. TURSI: We believe the risk management  
6 plan will effectively mitigate that risk, as I've  
7 described it. In terms of the first point and the  
8 specific physicians ultimately receiving access, we  
9 ultimately went to them to ask what do you prefer.  
10 Based upon your knowledge of the procedure, based upon  
11 your understanding of the disease, what would be the  
12 best method ultimately to provide training. And that  
13 answer came back overwhelmingly, not just from  
14 investigators but also generalists -- and when I say  
15 "generalists," meaning general rheumatologists,  
16 general surgeons within the specialties I told you.  
17 And that was the feedback we ultimately received.

18 In terms of the learning curve or the  
19 training curve, I would ask Dr. Kaplan or Dr. Waller  
20 to come up to speak specific to their example, because  
21 these were new physicians at using AA4500. They had  
22 not had access to this before. They had never used it



1 outside of their initial experience in the clinical  
2 program. So I think they could probably provide the  
3 best representation of what that "training," looks  
4 like.

5 Dr. Kaplan.

6 DR. T. KAPLAN: Sure. I think the most  
7 important part, honestly, of the training is to  
8 highlight the problem, which is tendon rupture. So we  
9 have to impart upon the physician they need to be  
10 concerned. They need to pay attention. They need to  
11 be surfing their e-mail if they're going to do a new  
12 procedure that they're just learning. As a surgeon,  
13 my training, you do it as during a residency. You  
14 learn procedures, but even after that, there's always  
15 new products, new techniques that are being developed.

16 And as a practicing physician, as you know,  
17 most of the time, that's not done in a hands-on  
18 workshop, per se. You have the experience that you  
19 have through your practice, through your training, and  
20 then you can adapt to new tools to your training.  
21 This is just one more tool that we've utilized.

22 I had no experience with collagenase prior

1 to my involvement in the trial. I will say that the  
2 first time I did it was -- again, I was kind of  
3 comfortable with the injection, but feeling that  
4 resistance of injecting into the cord was a new  
5 experience. But you knew it right away. It didn't  
6 take -- as you did that injection, you had a sense of  
7 what that injection was. And if you weren't in that  
8 cord and you lost resistance on your plunger, you knew  
9 immediately that potentially, you were out of that  
10 cord and you needed to stop that injection.

11           So I think the most important thing to  
12 highlight to anyone who's going to do this -- and I do  
13 agree with Auxilium that we should limit it to  
14 physicians who do understand the anatomy of the hand,  
15 and particularly the anatomy of Dupuytren's disease,  
16 because those cords can vary in patient to patient.

17           So we want to make sure we get physicians  
18 who are knowledgeable with the condition and who are  
19 going to adopt it and utilize a new treatment, and  
20 give it the due that it requires to learn it properly.

21           DR. DELCONTE: And Dr. Waller can also  
22 comment on the learning curve, if he could come

1 up -- because he's done a number of injections.

2 DR. WALLER: The one question you addressed,  
3 I think, or one answer, the tendon ruptures did not  
4 occur with any of the rheumatologists doing the  
5 injections. As Dr. Kaplan said, the first injection,  
6 yeah, it was a little -- it's certainly different, and  
7 subsequently, it was much more comfortable after that.  
8 I do think Auxilium's doing the best they can for a  
9 video.

10 And as other rheumatologists, I remember  
11 when we got our first dose of -- one of our biologic  
12 drugs that may be intravenous, we usually weren't set  
13 up back 10 years ago to have IV poles, and now we all  
14 have essentially, epinephrine, cortisone for allergic  
15 reactions. And unfortunately, there were no videos to  
16 watch a patient get some of these biologic drugs for  
17 us.

18 So to me, this video is actually again  
19 more -- making me more comfortable, and I think other  
20 rheumatologists would -- in the sense that it's a  
21 potent drug, certainly, but we deal with potent drugs  
22 every day. And, again, no video, no follow-up.

1 Certainly, any of the side effects are based on  
2 physicians calling in and making the description or  
3 the complaint, if you will, in the sense of what  
4 happened. And I think Auxilium's got it set up  
5 correctly.

6 DR. DELCONTE: And just regarding the  
7 question you had about the timing of the tendon  
8 rupture with regard to experience, there was no  
9 correlation to that. One of the three occurred in one  
10 of the investigators who was also a Phase 2  
11 investigator. And the numbers were really too small  
12 to look at other factors that could correlate with  
13 that.

14 DR. O'NEIL: Thank you. The next question  
15 is from Dr. Haque, and then Dr. Buckley.

16 DR. HAQUE: Thank you. I actually have  
17 several questions, so if I could, what I'll do is I'll  
18 just ask one now and then if Dr. O'Neil could indulge  
19 me later.

20 This question is directed towards Dr. Tursi  
21 regarding risk management. And it's regarding the  
22 patient education. Since this is sort of a new type

1 of procedure that we're going to be doing in the  
2 office, I was wondering what your thoughts are  
3 regarding creating a standardized consent form, and  
4 having that basically enumerate and list very  
5 specifically risks and benefits, and having all users  
6 provide that to their patients in getting informed  
7 consent so that it's not like just giving trigger-  
8 finger injections where people are very widely  
9 variable in how they approach that with their patients  
10 regarding risks and benefits.

11 DR. DELCONTE: The informed consent has not  
12 been part of the risk management program at this  
13 point. It's really the extensive patient information,  
14 the patient information leaflet and additional  
15 information. So we have not included that as part of  
16 the program yet.

17 DR. O'NEIL: Okay. Dr. Buckley.

18 DR. BUCKLEY: I think we're all trying to  
19 understand -- I guess the major concern is about  
20 tendon rupture, so I'm trying to understand why does  
21 that happen. Does it happen because the needle is put  
22 in the wrong place, or even if the needle is put in

1 the right place, can there be some extravasation that  
2 then leads to tendon rupture? And in that same line  
3 of questioning, I think all of us who have experience  
4 doing corticosteroid injections in hands know that  
5 sometimes there's tracking of the corticosteroid back  
6 through the skin.

7 Do you have much experience with what  
8 happens when there is tracking of this, or have you  
9 tried in animals to specifically put it in to a dermal  
10 area and see what reactions are?

11 And I have one other question after that.

12 DR. DELCONTE: Well, I'll let Dr. Tursi deal  
13 with the question about the tendon rupture, and then  
14 Dr. Hart can talk about what we've done, because we've  
15 actually misinjected deliberately this into a number  
16 of structures, so we can tell you what happens with  
17 that.

18 DR. TURSI: As to the specifics of the  
19 tendon rupture, there's no way to determine exactly  
20 what happened in terms of causing that rupture. We  
21 clearly attribute it to AA4500. Whether it was  
22 directly injected into the tendon or if it was

1     injected in the proximity of the tendon is unknown  
2     based on the specifics of the procedure.

3             So, again, although those numbers were  
4     small, it was something that was very important to us,  
5     and clearly is a key focus of our risk management  
6     plan.

7             I will ask Dr. Hart to come up and speak a  
8     little bit about the non-clinical work that you had  
9     asked about.

10            DR. HART: I'm going to point you to the  
11     results of two of our non-clinical studies, one of  
12     which will address your question on extravasation, and  
13     the other which will address your question of  
14     misplacement of the injection. The results are  
15     similar in both studies.

16            To address extravasation, I'm going to refer  
17     you to the first of these studies, which was our rat  
18     intravenous toxicity study. It was clear from having  
19     observed the injection sites histologically that in a  
20     few of these animals, there was some extravasation  
21     from the IV injection site. And in this location  
22     which is the rat tail, the injected veins are in very

1 close proximity to the skeletal muscle, the tendons,  
2 the arteries and the bones of the tail. So we  
3 basically had all of the structures represented that  
4 you'd see in a finger.

5           No effects on the injected vessel itself.  
6 Where it had extravasated, there were no effects on  
7 the tendon fibers directly, although the peritendon  
8 had lysed in some of the higher-dosed animals. The  
9 nerves, the arteries, the skeletal muscle, the bone  
10 and the collagen were all histologically normal. And  
11 when those tendons that had had the peritendon's lysis  
12 were evaluated two weeks later, there was evidence  
13 that that change was reversing.

14           And to answer your question about  
15 inadvertent administration, missing the cord and  
16 putting it into a subcutaneous location, I'll refer  
17 you to a series of three studies that were performed  
18 to support a different indication but will answer your  
19 question in terms of Dupuytren's disease, because the  
20 location is very similar. It was submucosal in the  
21 penis adjacent to the vein-artery nerve complex, as  
22 well as in different places. When the material was



1 injected submucosally or into the adventitia of the  
2 penis and overlaying the tunica albuginea, which is a  
3 dense collagen structure similar to a tendon, there  
4 was no evidence that leakage went down into the tunica  
5 albuginea and caused any lysis.

6           We did see the same sort of effects that  
7 were seen in the clinic, red blood cells and swelling,  
8 but no effects on arteries, on nerves and on larger  
9 veins. Only the smaller venules were disrupted. There  
10 were some changes in the walls of the arteries, some  
11 collection of red blood cells that was not associated  
12 with any damage to the smooth muscle, or  
13 interestingly, to the periarterial collagen. And that  
14 was verified by using a special stain, trichrome,  
15 which highlights collagen and collagen damage.

16           And, again, I want to point out that all of  
17 these effects reversed following withdrawal of the  
18 compound. There were no permanent effects in those  
19 arteries, even where this red blood cell accumulation  
20 occurred.

21           So we've evaluated extravasation. We've  
22 evaluated direct misadministration. And in all cases,

1 normal structures were spared, and in all cases, the  
2 changes reversed within two to four weeks following  
3 administration.

4 DR. BUCKLEY: And do you have any specific  
5 intradermal injections? You have extra -- but have  
6 you actually looked where you specifically put it  
7 intradermally?

8 DR. HART: Intradermally?

9 DR. BUCKLEY: Yeah.

10 DR. HART: There was a study done in support  
11 of that by the originator company. There was no  
12 histologic evaluation done, unfortunately. There's  
13 subdermal injections that were published in the  
14 literature. And, again, the same spectrum of changes  
15 is described, which is the inflammation, the bleeding  
16 and the reversibility of the effects. But those  
17 investigators didn't specifically talk about blood  
18 vessels and nerves.

19 I can tell you from the dog study that there  
20 was no upstream effects. In other words, the  
21 overlying mucosal cells and the interaction between  
22 the skin and the basement membrane were histologically

1 normal.

2 DR. T. KAPLAN: I was going to take an  
3 opportunity to kind of share with you clinically what  
4 we experienced. When tendon ruptures start -- before  
5 this multi-center Phase 3 trial, no tendon ruptures  
6 had occurred with the use of collagenase. So shortly  
7 after -- I don't know exactly how many months, but  
8 within the first several months after the study  
9 started, that two tendon ruptures occurred. And in  
10 response to that, we kind of as investigators got  
11 together to try to figure out was there any kind of  
12 pattern, is there anything that may be putting it more  
13 at risk?

14 There have only been two out of 1,000  
15 patients, so it's hard to draw conclusions. But those  
16 first two were both in the small fingers when treating  
17 PIP joints. And we know that as that cord kind of  
18 extends out toward that digit, as that cord gets  
19 closer and closer to the PIP joint, it gets closer and  
20 closer to where the flexor tendon system is. So  
21 certainly, it would affect -- with the way this drug  
22 works, that if it does get close to the tendon system,

1    then it could potentially cause risk and weakening of  
2    that system.

3                So we as investigators got together and we  
4    kind of went through the injection technique, made  
5    some clarifications to kind of tell investigators,  
6    hey, we should really stay away from into the finger  
7    area. And when treating a PIP cord, really target it  
8    near the base of the finger.

9                This is just kind of -- this is a slide that  
10   Dr. Tursi had shown of kind of that distance between  
11   the cord and that flexor tendon system. And, again,  
12   in the small finger which is not seen here,  
13   oftentimes, there's a central cord that comes down the  
14   center of the palm and goes right down the midline of  
15   the digit, which is clearly accessible and oftentimes  
16   will separate fairly far from that flexor tendon  
17   system.

18               In the small finger, oftentimes, there's  
19   something called an abductor digiti minimi cord, which  
20   is along the side of the digit. Some patients will  
21   actually have both of these cords, which will then  
22   kind of form a confluence as it goes over the --

1 around the PIP joint and just proximal to that.

2           So those are areas we felt that the  
3 injection, you have to be a little more careful or  
4 move that injection away from those areas to keep it  
5 away from the flexor sheath.

6           And then in the second question, as far as  
7 extravasation out of the skin, I certainly experienced  
8 that when I was doing it. I was much more happy. I  
9 definitely didn't want to extravasate deep to the  
10 cord, and some material would come up out of the skin.  
11 The only kind of side effects I saw from that, some  
12 patients did have some formation of blood blistering  
13 in the skin. That could have been due to the swelling  
14 that, we see that with fracture blisters. So it could  
15 be related to the swelling.

16           But potentially the collagenase, is hard to  
17 know. But when extravasated, usually, that cord is so  
18 close to the skin, you could actually see it leaking  
19 right through the skin.

20           DR. BUCKLEY: And just as a follow-up  
21 question to all this, it's clear with some experience  
22 and good understanding of the anatomy, there's a

1 learning curve. But for those who are less  
2 experienced, have you thought about things like  
3 ultrasound guidance?

4 DR. T. KAPLAN: I think that was actually  
5 done in some of the earlier Phase 2 trials, that you  
6 looked at ultrasound to map out the cord, to look at  
7 the distance between the cord and the tendon sheath  
8 underneath of it. Honestly, when you see patients  
9 with Dupuytren's disease, the cord is just right  
10 there. It's right underneath the skin, and it's hard  
11 to miss. So it's very easy to identify the cord and to  
12 get the injection to the cord. The key is not getting  
13 through the cord and putting the injection deeper to  
14 that or from the side.

15 So I personally don't feel that ultrasound  
16 would be all that beneficial in giving me better  
17 definition of the cord and where it is, because I  
18 think it's palpable.

19 DR. O'NEIL: Thank you. The next question  
20 is -- we'll go back to Dr. Haque and then Dr. Swartz  
21 and Ms. Aronson.

22 DR. HAQUE: I had a question regarding the

1    basic science that was presented earlier on, I think,  
2    Slide 9 regarding the collagenase types.  So am I  
3    correct that the Class 1 and Class 2 collagenase don't  
4    have anything to do with Type 1 versus Type 2  
5    collagen?  They're just separated by where they cleave  
6    the collagen fibers, and what types of collagen do  
7    they work on?  Have you seen any injuries to joint  
8    surfaces or other structures as well?

9               DR. DELCONTE:  To answer your question, I'll  
10   have Dr. Hart talk about the types of -- that is  
11   different than the types of collagen, and the types of  
12   collagen that the AA4500 has a preference for, a  
13   selectivity is Types 1 and Type 3.  And Dr. Hart can  
14   describe that a bit more.

15              DR. HART:  Your question about the  
16   collagenase classes relates to the -- there are two  
17   different forms of the enzyme that are secreted by the  
18   bacterium.  Each is a separate gene product.  They're  
19   a little bit different structurally, but they don't  
20   determine the substrate specificity.  Either one has  
21   the same substrate preferences.  They're active in a  
22   test tube against a wide variety of collagen subtypes,

1 but in vivo, it appears that their activity is  
2 somewhat selective for the fibrillar collagens, which  
3 is Type 1 and Type 3, with sparing of globular  
4 collagens such as Type 4, Type 6 and Type 8. And  
5 that, I think, translates to the effects we saw in the  
6 animal studies where there was no degradation of the  
7 periarterial collagen, which is primarily Type 4.

8 And if you had a second question, could you  
9 please repeat it?

10 DR. HAQUE: I think that was essentially it.

11 DR. HART: Thank you.

12 DR. O'NEIL: The next question is from  
13 Dr. Swartz, and just because he will be speaking to  
14 someone behind him, I'm going to remind him to speak  
15 into the microphone.

16 DR. SWARTZ: Thank you. I have two  
17 questions and a comment. First, most patients who  
18 come to my office with this condition have it in a  
19 mild form. They may have a nodule that may or may not  
20 be painful. They may have an early contracture. And  
21 our advice to these patients is that we don't know if  
22 it's going to be progressive or not. And so



1 observation, as has been mentioned earlier, is the  
2 most often the first encounter and the first advice to  
3 these patients, and they come back when it's more  
4 significant.

5           But with this medication, I can envision  
6 that our inclination is going to be to recommend that  
7 we treat them without knowing that in fact, they will  
8 have a progressive condition, and treat them before  
9 the contracture of the MP joint is more than 30  
10 degrees or the PIP joint more than 20 degrees. So my  
11 question to the FDA panel as well as to the Auxilium  
12 people is would this be considered an off-label  
13 treatment, and is this going to be -- and I guess, a  
14 better question, will there be a long-term focus and  
15 study of these patients to see if in fact, it does  
16 prevent progressive disease? That's my first  
17 question.

18           And then the second is, we haven't heard too  
19 much yet about the PIP joint contracture. On the  
20 opposite side of the spectrum is a severe contracture  
21 of the PIP joint to 70 or 90 degrees. And what has  
22 been the effectiveness of the injection in the PIP

1 joint patients to relieve that degree of contracture?  
2 Because I think this is where the most trouble is  
3 going to be seen. It's pretty straightforward, I  
4 think, to inject the palmar cord in the mid palm, but  
5 trying to relieve that PIP joint contracture where the  
6 spiral cord goes around the digital nerve and where  
7 you have a confluence, not only in the little finger  
8 but in the ring finger as well, of multiple abnormal  
9 structures surrounding the flexor tendon.

10 So we may want to see a stratification of  
11 patients and who's going to treat them based on the  
12 degree of severity, particularly in the PIP joint.

13 DR. DELCONTE: Let me address that second  
14 question first about the differences in joint and  
15 severity. We did a sub-analysis, and if we had the  
16 slide up, we can show you that in the two large  
17 multi-center studies, this is the responder rate here,  
18 the proportion of patients, and these are the four  
19 different subtypes.

20 And what you see first of all that is in the  
21 left two columns, the MP joints generally do better  
22 than the PIP joints. And joints generally of low

1 severity tend to do better than those of high  
2 severity. So in the high severe -- and we only  
3 stratified this. We sort of broke it in half, less  
4 than or equal to 40 and greater than 40. Here, about  
5 a quarter of the patients will achieve this zero to  
6 five degrees.

7           So this is what we see in the pooled  
8 studies. And when we were talking to and looking at  
9 the literature in hand surgery, it is that PIP joints  
10 generally as particularly the ones of high severity do  
11 not tend to correct as well.

12           Then furthermore, to answer the question  
13 about the labeling and where this would be used, as  
14 you saw, the clinical trials used a less than -- a  
15 contracture that was greater or equal to 20 degrees.  
16 And we would not be seeking an indication specifically  
17 for nodules. We would be only where there's a  
18 contracture and in most cases, the patients wouldn't  
19 be coming in unless they had some functional  
20 disability as well.

21           Regarding long-term follow-up, we do propose  
22 looking at a two- to five-year follow-up of not only

1 joints that have been treated, but joints that have  
2 not received therapy, to look for a progression. So  
3 we'll gain some additional information about the  
4 natural history of the disease. And what we  
5 understand from the literature is about half the  
6 patients with an early contracture or nodule will  
7 ultimately go on and progress. But this will give us  
8 some additional information on durability, on overall  
9 progression in untreated joints, and some additional  
10 long-term safety data.

11 DR. O'NEIL: Our next question is from  
12 Ms. Aronson.

13 MS. ARONSON: I'd like to start with an  
14 appreciation of the presentation. I found it very  
15 helpful as well as the briefing document. I also  
16 thought the video was a wonderful tool that could be  
17 used for continuing reference as physicians learn to  
18 use the product.

19 I was left with one question, and that is if  
20 there is a slide about exclusion of patient  
21 population. I know that Dr. Tursi talked about patient  
22 population, and Dr. DelConte referenced drugs such as

1 tetracycline and anticoagulants that were omitted.  
2 But he also said "other drugs," and I'm wondering what  
3 those other drugs are, and if they coordinate with the  
4 patient population that might have been omitted from  
5 the trial.

6 DR. DELCONTE: Let me put this slide up on  
7 our exclusion criteria that you referred to. And  
8 there were really two classes. Tetracycline,  
9 antibiotics were excluded because of a theoretical  
10 concern about inactivation of the collagenase. And  
11 this was just historically carried out through the  
12 studies.

13 The second class of drugs were  
14 anticoagulants, and this was because we know you could  
15 get some bleeding and bruising at the site. Other  
16 than we did allow low dose aspirin, but if a patient  
17 was anticoagulated, they were not allowed to be in the  
18 trial. And that would be also reflected in the label,  
19 and how we would suggest this be used.

20 MS. ARONSON: Patients with rheumatoid  
21 arthritis, for instance, would also be included in the  
22 trial?

1 DR. DELCONTE: In the clinical trials, we  
2 didn't want any types of illnesses that would confound  
3 measurement. So if they had any appreciable deformity  
4 or contractures of their fingers, we did exclude that  
5 patient population so that we'd be able to identify  
6 just the impact of the drug and not have any  
7 confounding from other diseases. So they were not  
8 included.

9 DR. O'NEIL: Dr. Mazor.

10 DR. MAZOR: This is a bit of a follow-up on  
11 Dr. Haque's question. And it relates to informing  
12 patients of the risks and benefits of the procedure.  
13 And you've talked some about the patient information  
14 packet, or however you refer to that. I'm wondering  
15 when that would be given to the patient, because I  
16 think there's a difference when you get something,  
17 look at this and stick your hand out kind of thing  
18 versus look at this, think about it and come back and  
19 tell me in a week or whatever amount of time.

20 DR. DELCONTE: Dr. Tursi can address the  
21 informed consent.

22 DR. TURSI: Ultimately, that would be at the

1   discretion of the individual physician, but as having  
2   been a physician in practice, I agree with you.  
3   Clearly, there is an advantage to providing patients  
4   with this information well in advance of any proposed  
5   procedure. So clearly, what we're trying to do as  
6   part of our overall risk management plan is not just  
7   rely on that patient product information leaflet, but  
8   also provide information to patients via website and  
9   other patient information brochures that would be  
10   available in physician offices. So they could gather  
11   that information, have a chance to digest in advance  
12   of the procedure.

13           DR. MAZOR: So I'm wondering -- and this  
14   kind of fits with the physician packets as well,  
15   because one could envision that the physician training  
16   materials or the physician attestation or commitment  
17   could include a commitment to informing patients in  
18   this way.

19           And related to that, I wondered about,  
20   there's kind of one way to find out if I know  
21   something and you ask me, and I can say yes even if I  
22   don't, and there's another way, which is you have some

1 level of testing me. You ask me some simple questions  
2 about do you know where to report an adverse event,  
3 kind of how can you find this information. Do you  
4 know when we recommend that you give this information?  
5 So like a lot of continuing medical education, some  
6 very straightforward questions that might be a part of  
7 that attestation.

8 DR. TURSI: Yes, we share your concern, and  
9 we absolutely appreciate your advice in that regard.  
10 I can show you the part of the draft attestation that  
11 I think directly addresses your question. At the  
12 bottom, we specifically ask physicians, "I will  
13 counsel each patient on the risks and benefits of  
14 AA4500 and provide each patient with the patient  
15 package insert."

16 So clearly, we are familiar with that issue,  
17 and we clearly want to provide as much information as  
18 possible, not just to physicians but to patients as  
19 well.

20 DR. O'NEIL: Thank you. We have now reached  
21 the witching hour, and we will take a short 10-minute  
22 break. Committee members, I'd like to remind you that



1   there should be no discussion of the meeting topic  
2   during the break among yourselves or with any member  
3   of the audience.

4                   And we will resume promptly at 10:45.

5                   (Whereupon, a recess is taken.)

6                   DR. O'NEIL: Now, I'd like to call on  
7   Dr. Eric Brodsky, who is a clinical reviewer at DAARP  
8   at the FDA, who will begin the FDA presentation.

9                   DR. BRODSKY: Good morning, Advisory  
10   Committee members. Good morning, members of Auxilium.  
11   Thank you for coming. My name is Eric Brodsky. I'm a  
12   medical officer at the FDA. The FDA appreciates your  
13   time and your efforts in helping us, advise us, about  
14   Auxilium's proposed application for Xiaflex, with the  
15   established name of collagenase clostridium  
16   histolyticum, for the proposed indication of advanced  
17   Dupuytren's disease.

18                   During my presentation, I will discuss the  
19   major efficacy and safety results of the application;  
20   I will highlight investigator training in the clinical  
21   trials and the proposed training of clinicians if  
22   Xiaflex were approved. And I will also provide a

1 benefit/risk assessment based upon the clinical trial  
2 data.

3           Auxilium presented a detailed background  
4 regarding Dupuytren's contracture. Auxilium also  
5 presented a detailed background of Xiaflex. Thus, I  
6 will only add that in 1996, Xiaflex was granted an  
7 orphan designation for the treatment of advanced  
8 Dupuytren's disease. In general, products can be  
9 given an orphan designation for specific indication if  
10 the disease will likely affect fewer than 200,000  
11 patients in the United States.

12           I will also emphasize that Auxilium proposes  
13 that Xiaflex be given by a physician experienced in  
14 the diagnosis and management of Dupuytren's disease.

15           There were six randomized double-blind  
16 placebo-controlled trials of Xiaflex in patients with  
17 Dupuytren's contracture. The only difference between  
18 our assessment and Auxilium's assessment of these  
19 trials is that we believe the two largest trials, with  
20 many sites and many investigators, served as the  
21 primary supports for the efficacy and safety of  
22 Xiaflex in Dupuytren's contracture. These trials are

1 Studies AUX-CC-857 and AUX-CC-859, abbreviated here as  
2 Studies 57 and 59 respectively.

3 Study 57 had a total of 308 treated patients  
4 at 16 U.S. sites. Study 59 had a total of 66 treated  
5 patients at five Australian sites. In these trials,  
6 patients must have had a fixed flexion contracture of  
7 at least 20 degrees of an MP joint or a PIP joint  
8 caused by a palpable cord to be included. Patients  
9 may have received up to three injections of Xiaflex or  
10 placebo directly into one cord given at four-week  
11 intervals. If the contracture persisted 24 hours  
12 after the injection procedure, the investigator  
13 extended the treated finger in an attempt to rupture  
14 the cord. Additional support for the  
15 efficacy and safety of Xiaflex in the treatment of  
16 Dupuytren's contracture comes from four smaller  
17 randomized double-blind placebo-controlled trials,  
18 abbreviated as Studies 02, 03, 51 and 53.

19 I will focus the efficacy presentation on  
20 the results from the two trials that served as the  
21 primary support for the efficacy of Xiaflex in the  
22 treatment of Dupuytren's contracture. The primary

1 efficacy endpoint for Studies 57 and 59 was the  
2 proportion of patients that achieved a reduction of  
3 the contracture of the primary joint, MP or PIP joint,  
4 to zero to 5 degrees 30 days after the last injection,  
5 where up to three injections could have been given.

6           Essentially, we are measuring the proportion  
7 of patients who achieve a straight joint, which is a  
8 clinically meaningful endpoint. In both trials, a  
9 statistically significantly greater proportion of  
10 Xiaflex-treated patients compared to placebo-treated  
11 patients achieved the primary efficacy endpoint after  
12 up to three injections.

13           In Study 57, the U.S. study, 64 percent of  
14 Xiaflex-treated patients, compared to 7 percent of  
15 placebo-treated patients, achieved the primary  
16 efficacy endpoint. In Study 59, 44 percent of  
17 Xiaflex-treated patients, compared to 5 percent of  
18 placebo-treated patients, achieved the primary  
19 efficacy endpoint.

20           For the Xiaflex-treated patients, the mean  
21 number of injections required for clinical success was  
22 1.7 in the two trials. It's important to note that

1 the proportion of Xiaflex-treated patients who  
2 achieved clinical success after the first injection  
3 was 39 percent in Study 57 and 27 percent in Study 59.

4           After up to three injections, Xiaflex  
5 treatment resulted in a greater mean decrease in the  
6 mean percentage change from baseline in the  
7 contracture of the primary joint. In Study 57, the  
8 baseline contracture was about 50 degrees. After  
9 Xiaflex treatment, the contracture was about 12  
10 degrees, resulting in a 79 percent reduction in  
11 contracture degree. In contrast in Study 57, placebo-  
12 treated patients demonstrated a 9 percent in  
13 contracture reduction. The results from the  
14 Australian study, Study 59, were similar to the U.S.  
15 study for this endpoint.

16           This is a representation of the efficacy of  
17 Xiaflex in the treatment of Dupuytren's contracture.  
18 These results are based upon the results from Study  
19 57. The white line represents a normal situation,  
20 where patients could extend their finger completely  
21 without a contracture, zero degrees of contracture.  
22 The yellow line represents the mean baseline severity

1 of contracture in Study 57, which was about 50  
2 degrees. This is prior to the injection. The green  
3 line represents the mean degree of contracture after  
4 up to three Xiaflex injections, which is about 12  
5 degrees. For Study 57, the contracture was reduced  
6 close to normal after Xiaflex treatment.

7           Contracture recurrence is a concern for any  
8 treatment for Dupuytren's disease because of the  
9 nature of the disease, which is progressive and  
10 incurable. Few Xiaflex-treated patients in the  
11 studies experienced a recurrence, approximately 4  
12 percent. However, the follow-up period was very  
13 limited. The mean time of follow-up was about seven  
14 months. In the Xiaflex studies, recurrence was  
15 defined as a return of the contracture greater or  
16 equal to 20 degrees associated with the presence of a  
17 palpable cord in patients who initially experience  
18 clinical success.

19           In an attempt to provide some perspective on  
20 the incidence of recurrence following Xiaflex  
21 treatment, we looked at the published literature for  
22 the incidence of recurrence from the most common types

1 of surgery for Dupuytren's contracture: fasciotomy and  
2 fasciectomy. Fasciotomy, as mentioned before, is a  
3 division of the cord and is usually done  
4 percutaneously. Fasciectomy is a more-extensive  
5 procedure, in which the diseased fascia and sometimes  
6 the normal surrounding fascia are removed.

7           Using a more-severe definition of  
8 recurrence, severe enough to require another surgery,  
9 with a much longer duration of follow-up, ranging from  
10 two years to about 10 years, we found a wide range of  
11 recurrences after surgery. The incidence of  
12 recurrence ranged from zero percent following  
13 dermofasciectomy, which is a more extensive form or  
14 type of fasciectomy, to up to 66 percent following  
15 fasciotomy.

16           One concern is that physicians with  
17 different expertise may have different efficacy  
18 results. To shed some light on this issue, we  
19 performed an exploratory subgroup analysis using the  
20 primary efficacy endpoint by expertise of the  
21 investigator who performed the injections. In pooled  
22 Studies 57 and 59, the majority of the injections were

1 performed by hand surgeons. Approximately 81 percent  
2 of the injections were performed by hand surgeons,  
3 whereas about 15 percent of the patients were injected  
4 by orthopedic surgeons, and about 4 percent of the  
5 patients were injected by rheumatologists.

6           Within each study, investigators  
7 irrespective of specialty obtained similar results for  
8 the primary efficacy endpoint, as you can see here.  
9 Although there were no major differences in efficacy  
10 results for each of the specialty groups, no  
11 definitive conclusions can be drawn because of the  
12 limited number of patients who were injected by non-  
13 hand surgeons.

14           Now let's turn our attention to the safety  
15 assessment. There were two populations used for this  
16 safety analysis. First were patients in the  
17 randomized double-blind placebo-controlled portions of  
18 pooled Studies 57 and 59 through Day 90. In this  
19 pooled safety database, about 250 patients were  
20 treated with Xiaflex, and 125 patients were treated  
21 with placebo. The Xiaflex dose was .058 milligrams.

22           The safety of Xiaflex was also evaluated in



1 the controlled and uncontrolled portions of all 12  
2 submitted Xiaflex studies. In this pooled safety  
3 database, about 1100 patients were treated with  
4 Xiaflex, representing about 2600 injections. The mean  
5 duration of safety follow up for these patients was  
6 about 10 months. About 60 percent of patients  
7 received two or more Xiaflex injections. You can see  
8 the distribution of Xiaflex injections within this  
9 table.

10 We analyzed the safety of Xiaflex in the  
11 controlled portions of the pooled Studies 57 and 59  
12 who received up to three injections of study  
13 medication. No one died in the controlled period.  
14 There was a slightly greater proportion of Xiaflex-  
15 treated patients compared to placebo-treated patients  
16 who had a serious adverse event. This difference was  
17 entirely due to serious adverse events of the injected  
18 extremity.

19 A slightly greater proportion of patients  
20 had an adverse event leading to discontinuation, or a  
21 DAE. Two of the three patients in the Xiaflex group  
22 who had an adverse event leading to discontinuation,

1 the adverse event involved the injected extremity.

2 Almost all of the Xiaflex-treated patients had an

3 adverse event. The Xiaflex group had two times the

4 number of adverse events compared to the placebo-

5 treated group, patients after up to three injections.

6 We also analyzed the major safety results in

7 the controlled and uncontrolled portions of the 12

8 submitted Xiaflex studies on a per-patient basis, the

9 upper portion of the table, and on a per-injection

10 basis, the lower part of the table. In the controlled

11 and uncontrolled portions of the studies, five

12 patients died. The causes of death in the Xiaflex

13 clinical program appear to be consistent with what

14 might be expected for the underlying patient

15 population.

16 Eleven Xiaflex-treated patients had a

17 serious adverse event of the injection extremity. Of

18 these 11 patients, as mentioned before, three had a

19 flexor tendon rupture, which were likely related to

20 Xiaflex treatment.

21 We evaluated all the deaths that occurred in

22 the 12 submitted studies and in the pilot academic

1 study. There were seven deaths. All these patients  
2 received a 0.58 milligram dose of Xiaflex. There were  
3 no deaths in a limited number of placebo-treated  
4 patients. The seven deaths in the Xiaflex group were  
5 not expected, given the background co-morbidities of  
6 these patients. There appeared to be no relationship  
7 between the number of Xiaflex injections and the  
8 incidence of death. Finally, most of the deaths  
9 occurred six months after the last Xiaflex injection.

10 All the serious adverse events of the  
11 injected extremity occurred in patients who received  
12 0.58 milligrams of Xiaflex. The upper part of the  
13 table shows the serious adverse events during the  
14 controlled portions of Studies 57 and 59 through  
15 Day 90, and the lower part of the table shows the  
16 serious adverse events of the injected extremity in  
17 the open-labeled uncontrolled portions of the Xiaflex  
18 studies.

19 Of the 11 serious adverse events shown,  
20 seven, or 64 percent, occurred within two weeks of the  
21 last injection. Many of these patients required  
22 surgery or other medical therapy to correct this

1 serious adverse event. Of these 11 serious adverse  
2 events, three were flexor tendon ruptures, as  
3 mentioned by the applicant. All of them occurred  
4 within seven days of the last injection.

5 All three tendon ruptures occurred after  
6 Xiaflex was injected into a cord affecting the PIP  
7 joint of the fifth digit. All the tendon ruptures  
8 were likely related to Xiaflex treatment.

9 Other serious adverse events of note  
10 included a pulley rupture, as mentioned before, and  
11 complex regional pain syndrome, as mentioned before.

12 To see if the frequency of the serious  
13 adverse events involving the injected extremity was in  
14 the same ballpark as surgical complications following  
15 surgery for Dupuytren's contracture, we performed a  
16 literature search of surgical complications following  
17 fasciectomy and fasciotomy. The incidence of  
18 intraoperative complications such as arterial injury  
19 or nerve injury was approximately zero to 10 percent,  
20 and the incidence of postoperative complications range  
21 from zero to 18 percent.

22 The incidence of serious adverse events of

1 the treated extremity observed in the Xiaflex studies  
2 did not appear out of proportion to the incidence of  
3 surgical complications as reported in the published  
4 literature.

5           After up to three injections, two times as  
6 many Xiaflex-treated patients than placebo-treated  
7 patients had an adverse event. The overwhelming  
8 majority of Xiaflex-associated adverse events were  
9 local reactions. The most commonly reported  
10 Xiaflex-associated adverse events were hand edema of  
11 the injected extremity, contusion, injection site  
12 hemorrhage and extremity pain. These events were  
13 likely related to Xiaflex injection. After one  
14 injection, 95 percent of Xiaflex patients had an  
15 adverse event.

16           Xiaflex contains foreign proteins, so  
17 allergic reactions would not be unexpected,  
18 particularly with repeated exposures. However, there  
19 were no severe reactions, including those associated  
20 with respiratory compromise, hypotension, or end-organ  
21 dysfunction.

22           We performed an exploratory analysis of

1 pruritus adverse events. Xiaflex-treated patients had  
2 a greater proportion of pruritus adverse events  
3 compared to placebo-treated patients in Studies 57 and  
4 59. The incidence of pruritus increased in the  
5 Xiaflex treatment group with more injections. Thus,  
6 there's some evidence of mild allergic reactions  
7 associated with Xiaflex injections. However, there  
8 were no severe allergic reactions.

9           As mentioned previously, Xiaflex contains  
10 foreign proteins. Therefore, we would expect to see  
11 antibodies to both components of Xiaflex, AUX-I and  
12 AUX-II. We looked at the frequency of these  
13 antibodies and evaluated if they had any clinical  
14 consequences. After the first injection,  
15 approximately 86 percent of patients had positive  
16 antibodies to AUX-I and/or AUX-II. After the fourth  
17 injection, all Xiaflex-treated patients had antibodies  
18 to AUX-I and AUX-II.

19           However, there appeared to be no effects of  
20 these antibodies on the efficacy or safety of Xiaflex.  
21 Patients who developed neutralizing antibodies to AUX-  
22 I or AUX-II had similar efficacy as patients with

1 neutralizing antibodies.

2           Now I'm going to talk about special  
3 considerations for this application. Since the  
4 clinical trial results were based on experienced  
5 investigators who were highly trained in Xiaflex  
6 injections, it is important to compare the training of  
7 the investigators in the clinical trials to the  
8 proposed training of clinicians if Xiaflex were  
9 approved.

10           As mentioned by Auxilium, no hands-on  
11 training such as simulations were performed in  
12 preparation for the trials, and no simulations are  
13 planned for clinicians in practice if Xiaflex is  
14 approved.

15           As mentioned by Auxilium, investigators in  
16 Studies 57 and 59 received training manuals and DVDs.  
17 Auxilium also proposes to provide clinicians with  
18 manuals and a narrated video, as you've seen.  
19 Investigators in the trials attended workshops and  
20 meetings regarding injection technique, although not  
21 all investigators participated, as you heard before.  
22 Instead of these type of workshops, Auxilium proposes

1 to provide personal liaisons to clinicians in  
2 practice.

3 In addition to the stated training for  
4 clinicians, Auxilium proposes additional risk  
5 minimization in the form of a managed distribution  
6 program that requires a physician to sign a form prior  
7 to receiving Xiaflex. Physicians must agree that they  
8 understand injection procedures and the risks of  
9 Xiaflex injection, including tendon rupture. If  
10 physicians do not sign the form, Xiaflex will not be  
11 provided.

12 Now I'm going to assess the benefits and  
13 risks of Xiaflex as seen in the clinical trials. The  
14 benefit-risk assessment of Xiaflex is based upon the  
15 pooled results of the controlled portions of Studies  
16 57 and 59 through Day 90, after up to three injections  
17 of study medication. These results may not be  
18 reflective of results in clinical practice.  
19 Nonetheless, this assessment may be useful as a  
20 starting point for your discussions.

21 Starting with the benefit of Xiaflex, note  
22 again all these benefits are based upon after up to



1 three injections. In the pooled clinical trials,  
2 again after up to three injections, two patients  
3 needed Xiaflex treatment to obtain the benefit of a  
4 straight joint in one patient. Also, after up to  
5 three injections, one patient needed Xiaflex treatment  
6 to obtain the more modest benefit of improvement of 50  
7 percent of the contracture degree in one patient.

8           Now moving on to the risks of Xiaflex, again  
9 note all the risks are also based on up to three  
10 injections of study medication in pooled Studies 57  
11 and 59. One patient needed Xiaflex to have one  
12 patient develop a local adverse reaction such as hand  
13 edema, contusion or pain of the extremity.

14           Now for the more serious events. 125  
15 patients needed Xiaflex for one patient to have a  
16 tendon rupture, and 83 patients needed Xiaflex  
17 treatment for one patient to have a serious adverse  
18 reaction other than a tendon rupture, such as complex  
19 regional pain syndrome or a pulley rupture.

20           In summary, results from the controlled  
21 trials demonstrate a statistically significant  
22 increase in the proportion of patients achieving

1 almost complete contracture reduction when treated  
2 with Xiaflex compared to placebo. Xiaflex injection  
3 was associated with twice as many adverse events  
4 compared to placebo, with most being local reactions.  
5 Serious adverse events including tendon ruptures were  
6 not common. Clinical trial results may represent a  
7 best case scenario, where the investigators had  
8 extensive professional training and were highly  
9 trained in Xiaflex injection and finger extension  
10 procedures.

11 Thank you.

12 DR. O'NEIL: Next, Dr. Kathryn O'Connell  
13 will speak to us about risk management considerations  
14 in the FDA approval process.

15 DR. O'CONNELL: Good morning. My name is  
16 Kathryn O'Connell. I'm with the Office of  
17 Surveillance and Epidemiology at FDA, the Division of  
18 Risk Management.

19 The FDA's concept of risk management is  
20 actually the overall and continuing process of  
21 minimizing risk throughout a product's life cycle to  
22 optimize the risk/benefit balance. And the reason

1     that the Division of Risk Management is here today is  
2     because there is a risk management issue that we've  
3     already talked about this morning, and that pertains  
4     to training, the role of training, and is required  
5     training necessary for safe use of this product.

6             It's an issue because the relationship  
7     between tendon rupture and improper administration of  
8     the product is unknown, and there's several factors  
9     that go into that. One is the generalizability of  
10    clinical practice of trial results that are obtained  
11    by highly trained investigators, and another issue is  
12    the unknown relationship for this product between  
13    tendon rupture and user factors such as specialty or  
14    hand anatomy expertise. And then there's the inherent  
15    potential damaging effect of the collagenase on  
16    collagen-containing structures adjacent to the cord.  
17    And your handouts should say the Dupuytren's cord.

18            In general, risk management for product  
19    safety issues are managed through the product's  
20    package insert, which all products have. Sometimes  
21    the sponsors provide extra education or training.  
22    Sometimes, there's post-marketing studies that are

1 involved, and there's always post-marketing  
2 surveillance.

3           However, if the seriousness of risk  
4 associated with any product or with this product  
5 specifically make it necessary to require and enforce  
6 training, then the Food and Drug Administration  
7 Amendments Act, or FDAAA, as you've probably heard it  
8 called, does provide FDA with the authority to require  
9 something called risk evaluation and mitigation  
10 strategies, or REMS. And accordingly, REMS can be  
11 required if and only if the FDA determines that these  
12 strategies are necessary to ensure that the benefits  
13 of the drug outweigh the risk.

14           REMS in general include one or more of the  
15 following: One is a patient medication guide. Second  
16 is a communication plan, and that's for healthcare  
17 professionals. And the last one is something called  
18 Elements to Assure Safe Use, and I'll talk more about  
19 that in a minute. But these often involve some form  
20 of restricted distribution. That may be how you've  
21 heard them referred to.

22           The first, the medication guide, this

1 provides for FDA-approved patient-friendly labeling,  
2 and it's required. The person who dispenses the  
3 product is required to give this to the patient. A  
4 patient medication guide can be required by the FDA if  
5 the FDA determines that one or more are true: First,  
6 is that patient labeling could help prevent a serious  
7 adverse event or events. The second is that the  
8 product has serious risks that could affect the  
9 patient's decision to use or continue to use the  
10 product. And the third is that patient adherence to  
11 directions would be crucial to product effectiveness.

12           The second thing I mentioned is called a  
13 communication plan. As I said, this is for healthcare  
14 providers. And a communication plan provides  
15 FDA-approved materials that are used to aid the  
16 sponsor's implementation of REMS, and/or inform  
17 healthcare providers about serious risk. And you're  
18 probably familiar with the "Dear Healthcare  
19 Professional" letters that you may have received about  
20 products.

21           These and other educational materials have  
22 been required in the past to alert prescribers to

1 serious risks associated with the use of certain drugs  
2 and biologics. But frankly, we don't know what the  
3 impact is of such letters.

4           The last thing that I mentioned as a  
5 component of REMS is something called Elements to  
6 Assure Safe Use. And there are six main categories of  
7 these elements, and I want to just note, because the  
8 sponsor had used the word "mandatory," but mandatory  
9 on this slide means, as I said on the previous slide,  
10 that the FDA would require and enforce, so that's the  
11 meaning of mandatory on these slides.

12           So the six items here are mandatory  
13 prescriber training or certification, mandatory  
14 certification of dispensers, drug administration  
15 restricted to certain healthcare settings -- for  
16 example, a hospital or an infusion center or  
17 whatever -- mandatory documentation of safe use prior  
18 to dispensing, mandatory monitoring of patients, and  
19 mandatory enrollments of patients in a registry.

20           As you can see from that list, Elements to  
21 Assure Safe Use are the three kinds of REMS that I  
22 talked about would provide the most strict control

1 over whether the product is used as per FDA-approved  
2 labeling. The downside is that these Elements to  
3 Assure Safe Use can impose significant burdens on the  
4 healthcare system and reduce patient access to  
5 treatment.

6 Therefore, Elements to Assure Safe Use  
7 should only be used if the product would otherwise not  
8 be approved due to specific serious risks listed in  
9 the labeling.

10 And in fact, the statute requires -- this is  
11 the wording out of the statute -- requires that  
12 Elements to Assure Safe Use must be commensurate with  
13 specific serious risks listed in the labeling. It  
14 cannot be unduly burdensome on patient access to the  
15 product and to the -- and they have to minimize the  
16 burden on the healthcare delivery system to the extent  
17 practicable, conform with elements for other drugs  
18 with similar serious risks, and be designed for  
19 compatibility with established distribution,  
20 procurement and dispensing systems for drugs.

21 So in summary, FDA does have the authority  
22 to require REMS if additional measures -- in this

1 case, required training -- are necessary to assure the  
2 benefits of CCH outweigh the risk. However, the risk  
3 management for CCH is for all products. It should  
4 minimize healthcare system burden and barriers to  
5 patient access to the extent possible within the risk  
6 mitigation goals.

7 Thank you.

8 DR. O'NEIL: Thank you. At this point, we  
9 will take some questions from the Committee to the  
10 representatives of the FDA. And the first one to  
11 raise his hand is Dr. Weisman.

12 DR. WEISMAN: Thank you, Kathleen.

13 I don't know whether Eric or Kathryn, which  
14 one should respond to this, but I'll just ask the  
15 question.

16 DR. O'NEIL: They're side by side.

17 DR. WEISMAN: You've pointed out in your  
18 presentation that the level of expertise and  
19 experience in doing these injections was limited  
20 almost exclusively to hand surgeons, and very few  
21 internists/rheumatologists were involved. And  
22 therefore, your presentation indicates that there was



1 not enough information to judge whether with this  
2 particular specialty or expertise of these clinicians  
3 was sufficient to allow the process to go forward  
4 safely.

5           Since we've heard from the sponsor that  
6 their process for screening individuals consists of  
7 filling out a form, and that form states just what  
8 your specialty is, and that includes rheumatologists,  
9 there's no scrutiny further as to additional expertise  
10 and then those people would automatically be allowed  
11 to use the -- to be able to use the procedure. Is  
12 there sufficient concern that you have that given the  
13 safety and the risk associated with this drug, that  
14 going forward, that this should be limited to hand  
15 surgeons as defined -- and we can ask for a moment  
16 what the definition of a hand surgeon is -- only and  
17 not opened up to generalists, internists or  
18 rheumatologists?

19           Just given the information we have so far on  
20 the safety and risk of this drug, is that what your  
21 concern is? And I'm trying to understand this.

22           DR. OKADA: That really is sort of the crux

1 of the issue that we're asking the Committee to advise  
2 us on, is just that we have these very nice study  
3 results and they're very -- and they're limited in  
4 terms of the background and the investigators. So how  
5 do we bring that forward to clinical practice? That's  
6 what we'd like you to comment on.

7 DR. WEISMAN: So that's sort of the crux of  
8 the matter here?

9 DR. OKADA: Uh-huh.

10 DR. O'NEIL: For the record, those comments  
11 were from Dr. Sarah Okada.

12 DR. WEISMAN: The other side of the question  
13 is, can we get a definition from our colleagues on the  
14 panel as to what constitutes a hand surgeon? I know  
15 from long experience and having distinguished over the  
16 years colleagues, associates of mine like Rich  
17 Gelberman and Dick Braun and Myles Cohen, I know what  
18 a hand surgeon is. But can you define for us what  
19 level of training and certification goes along with a  
20 hand surgeon, someone that might be, for instance,  
21 privileged at our institution to be able to do this  
22 procedure or similar procedures on Dupuytren's

1 patients?

2 DR. O'NEIL: I will recognize Dr. Kaplan for  
3 this.

4 DR. S. KAPLAN: Membership in the American  
5 Society for Surgery of the Hand requires that an  
6 individual be board-certified in either plastic  
7 surgery, general surgery or orthopedic surgery, and  
8 then has done a one-year fellowship in hand surgery.  
9 I think that's the current definition. Twenty years  
10 ago, there were many routes without certifying bodies,  
11 and there is no individual board certification in hand  
12 surgery. But there is something called a certificate  
13 of added qualification, which is administered by the  
14 boards of general surgery, orthopedic surgery and  
15 plastic surgery, which requires that one-year  
16 fellowship in hand surgery.

17 So I think that would be a definition, but  
18 I'd also like just comment, the procedure of needle  
19 aponeurotomy was developed by rheumatologists -- or  
20 popularized by rheumatologists in France. So I'm not  
21 sure we can -- we should exclude rheumatologists as a  
22 whole in this conversation.

1 DR. O'NEIL: The next question is from  
2 Dr. Saag.

3 DR. SAAG: I, like Dr. Weisman, share the  
4 concern that certain types of providers may have less  
5 experience. That's not to say that having a label as  
6 a hand surgeon or rheumatologist makes you distinctly  
7 qualified or unqualified to do this. But I do believe  
8 that a certain level of training and acquiring certain  
9 sufficient knowledge and skills is necessary to safely  
10 perform a procedure that has some risk.

11 And I want to ask the FDA about the specific  
12 mechanisms and perhaps examples about what might  
13 constitute mandatory prescriber training or  
14 certification. And beyond saying that it's mandatory,  
15 is there a way to assure that the training and  
16 certification leads to some measurable gain in  
17 knowledge or skills?

18 DR. O'CONNELL: That's a very good question.  
19 There are, as you know, REMS out there that have been  
20 approved that include the physician attestation or the  
21 healthcare provider attestation that they have the  
22 training needed to either understand the indication or

1 use the drug. I'm not really aware of any that  
2 actually measures that, like gives a test or they have  
3 to go to a hospital and show that they know how to do  
4 the procedure. I'm not sure. I'm not aware of any  
5 example like that. It's not to say that we couldn't  
6 try to design something like that, but right off the  
7 top of my head, I can't imagine what that would be.

8 DR. SAAG: For certain surgical procedures  
9 and devices, has the FDA required a practical training  
10 experience as part of the mandatory requirement?

11 DR. O'CONNELL: You mean for use of devices?

12 DR. SAAG: Say a new surgical procedure or  
13 device, yes.

14 DR. O'CONNELL: I'm not aware of any. Are  
15 you?

16 DR. RAPPAPORT: Unfortunately, we don't have  
17 anybody from that center. Those products are located  
18 in a separate center, and we don't regulate surgical  
19 procedures. So devices being in a separate center, I  
20 don't think there's anybody here who would know about  
21 that, but we can try to get that information for you.

22 DR. O'NEIL: Dr. Swartz is next.

1           DR. SWARTZ: I'd like to address the  
2 question of who are hand surgeons a little bit more  
3 broadly. I sit as the director of the American Board  
4 of Plastic Surgery, and have for the past six years  
5 been on the committee for training and certifying hand  
6 surgeons in this country. The American Board of  
7 Plastic Surgery and the American Board of Orthopedic  
8 Surgery and the American Board of General Surgery all  
9 have agreed that specific training in hand surgery and  
10 certification should follow the plan that was just  
11 described by Dr. Kaplan.

12           However, the facts of the matter on the  
13 ground are that there are many, many people who do  
14 hand surgery who are not board-certified or  
15 certificate of added qualifications in hand surgery  
16 physicians. And I'm not aware of very many hospitals  
17 that require that certificate to take hand call. If  
18 that were the case, we would have a woeful dearth of  
19 people able to treat hand patients on an emergency  
20 basis.

21           And for that reason, we don't have as a  
22 requirement, at least in our hospital in Pittsburgh,

1    that you be certified in hand surgery to treat hand  
2    patients.  You do need to have a certificate in  
3    general surgery or orthopedic surgery or plastic  
4    surgery and have experience with hand patients and  
5    demonstrate that experience in order to be accepted by  
6    the hospital for your privileges.

7                So this really comes down to privileging in  
8    a hospital setting for a surgical procedure, and in an  
9    outpatient clinical setting, there is no regulation.  
10   There is absolutely no regulation in this United  
11   States for the treatment of patients in an outpatient  
12   setting other than a surgical center facility, and  
13   that has to be kept in mind when we talk about who's  
14   going to treat these patients and what the  
15   risk/benefit ratio is.

16               My own personal opinion about this is that  
17   anyone who has experience treating hand patients and  
18   treats them regularly should be allowed to use this  
19   medication, and probably will use this medication.  
20   And a perfect example of that would be a rural  
21   physician, in the old style of the old general surgeon  
22   who sees all comers for all kinds of problems.  He'll

1 have the maturity and ability to decide about his risk  
2 profile and either will or won't use it based on that.  
3 And I think that's where this is going to come down  
4 to.

5 DR. O'NEIL: Dr. Haque has the next  
6 question.

7 DR. HAQUE: I'd like to second a lot of what  
8 Dr. Swartz said. The only thing in addition to what  
9 Dr. Kaplan was saying about actual certification for  
10 hand surgery is that it also requires submission of a  
11 case list with a broad base of experience in the prior  
12 year to taking an exam for certification that shows  
13 that you have experience in several different types of  
14 hand surgery, including tendon surgery, bone and joint  
15 or fracture surgery, microsurgery or perhaps  
16 congenital hand surgery.

17 So again, everybody's experience level is  
18 different, even within hand surgery, and I think what  
19 Dr. Swartz is saying is appropriate. I do think that  
20 a person who does several injections a month for  
21 trigger fingers would have the dexterity and the feel  
22 for how to give this injection, and I don't think that



1 we should necessarily exclude them based on some  
2 labeling that -- with their training or background.

3           The other point regarding devices, I don't  
4 know if it's FDA-mandated, but I have had experience  
5 with several devices, if I'm allowed to say,  
6 endoscopic carpal tunnel release and certain types of  
7 implant placements for joint replacements in the  
8 fingers where prior to getting approval to do it by  
9 the company, I actually had to do a hands-on course  
10 where I did cadaver training and listened to several  
11 lectures. Did not have a test, but actually had to  
12 perform the procedure and people were watching it.

13           I think that's obviously a huge additional  
14 burden on the provider and the company that's  
15 marketing the product, but in addition in this  
16 situation, it's a little bit hard to do. You can't  
17 exactly get cadavers that have lots of Dupuytren's and  
18 go in there and practice that. So actually getting a  
19 hands-on feel for this is going to be an on-the-job  
20 situation.

21           DR. O'NEIL: Dr. Buckley.

22           DR. BUCKLEY: I think what we're trying to

1 get a picture of here is what's the need in terms of  
2 patient need and what are our provider resources. So  
3 we want to make sure that the most experienced  
4 providers deliver this care, but on the other hand, we  
5 want to make sure that patients have access to the  
6 care. And I think that's where the dilemma is. We're  
7 calling it an orphan disease, which makes me think we  
8 don't really need to have a huge broad array of  
9 providers, although there will always be that patient  
10 in some remote area who might not have access.

11           It sounds like part of the answer might be  
12 that's it's providers who do a lot of hand work who  
13 have a lot of experience with this, so that we  
14 wouldn't want a provider who does an injection about  
15 once a year to be doing this kind of procedure. And  
16 that might be something else to take into  
17 consideration.

18           I think something we can't forget, I don't  
19 know what the reimbursement for this procedure is  
20 going to be. But I think we do know in clinical  
21 office-based practice that sometimes there's a bias to  
22 doing procedures by practitioners if the reimbursement

1 is high. And so that can lead to practitioners who  
2 maybe don't have a lot of experience doing the  
3 procedure maybe deciding this is something they should  
4 try to get more experience with or do more of. I  
5 think we have to be aware of that, that there might be  
6 some abuse of this procedure by people who not as  
7 experienced.

8           So it sounds like the challenge is what's  
9 the definition of an experienced person giving  
10 more-invasive hand care if not surgical care, but I  
11 think we have to be sensitive to the fact that there  
12 needs to be some restriction on this. And probably  
13 that shouldn't be up to the practitioner.

14           DR. O'NEIL: Dr. McAlindon.

15           DR. MCALINDON: Thank you. I'm trying to  
16 put the risk of tendon rupture into clinical context.  
17 Given that rheumatologists as well as hand surgeons  
18 inject complex small structures in hands and wrists  
19 rather regularly, I'm wondering if the FDA in their  
20 research found data to look at the overall risk, for  
21 example, of tendon rupture following peritendon  
22 corticosteroid injection, which is something we do

1 daily, and perhaps look to see if there are  
2 differences between rheumatologists and hand surgeons.

3           And the second part of this question, I'm  
4 wondering -- and the data from the clinical that show  
5 I realize are rather small in terms of that adverse  
6 event -- whether there was any signal, in fact, that  
7 the performer of the injection interacted with the  
8 level of risk. In other words, was there something  
9 about who did the injection that somehow mediated part  
10 of the risk or not? It's a two-part question.

11           DR. OKADA: This is Sarah Okada. In terms  
12 of your first question, we didn't actually perform a  
13 literature search to see what the going rate of tendon  
14 ruptures with peritendon steroid injections would be,  
15 but that's a useful suggestion and we'll take that  
16 back.

17           In terms of the details of who injected the  
18 patients who experienced the tendon ruptures, I  
19 believe -- and Eric, you can correct me if I'm wrong,  
20 but I believe that all of them, in fact, were injected  
21 by hand surgeons, which is sort of consistent with the  
22 fact that there were mostly hand surgeons doing the

1 procedures.

2 DR. O'NEIL: I'd like to ask a simple  
3 question, and this probably goes more to the sponsor.  
4 I know that's not quite right, but it's pertinent to  
5 the discussion at hand, which is was there any  
6 evidence that any of the different proposed methods of  
7 education had a bearing on either the success of the  
8 procedure or the ability to avoid complications?

9 DR. DELCONTE: We were not able to do the  
10 type of training that the investigator or  
11 sub-investigator had performed and the outcome, that  
12 just wasn't possible from the way we collected the  
13 data.

14 DR. TURSI: Just one comment, what I can  
15 also do, though, is provide a little context in  
16 regards to comparison of what the investigator  
17 training looked like versus what we're proposing, with  
18 your permission.

19 What I've done with this particular slide is  
20 I've kind of put side-by-side the injection training  
21 of the investigator versus the proposed physician  
22 training. And what ultimately we're proposing, we

1 believe is actually not only improved the investigator  
2 training, but includes additional facets to that  
3 training. When we consider the first area of  
4 training, which is the preparation of injection  
5 technique and finger extension, naturally, that was  
6 included as part of the investigator training. But  
7 we've gone ahead and improved that, and I can get into  
8 detail, should you desire, as opposed to going through  
9 the other points.

10           In terms of adverse event reporting,  
11 clearly, that would have been included as either part  
12 of the investigator brochure or part of the study  
13 protocols. But we've actually now encapsulated all  
14 that information in one structure, and that being the  
15 proposed training program. So physicians don't need  
16 to go to multiple places to get that information.  
17 They've got it all at their fingertips in one  
18 resource.

19           As it relates to important safety  
20 information and adverse event descriptions, again,  
21 we've gone beyond what we did in the investigator  
22 training. Other areas that weren't included in a

1 specific investigator training -- injection training  
2 specific to the joint, the risk of tendon damage,  
3 frequently asked questions, self-assessment,  
4 sequential completing of training being required prior  
5 to attestation and attestation being required before  
6 use -- they are all new additions to our proposed  
7 training program.

8           So I appreciate the opportunity to add that.  
9 Thank you.

10           DR. O'NEIL: Dr. Weisman, you have a  
11 question.

12           DR. WEISMAN: Yes, I want to -- question  
13 back to the FDA, though. I agree with Lenore. I  
14 think we need to focus and get away from this sort of  
15 food chain issue discussion, and talk about what  
16 really is the crux of the matter here, which is the  
17 discrepancy between what was done in the clinical  
18 trial and what's being proposed for safety monitoring  
19 and safety assurance in what the sponsor has given us.  
20 And now there's another discrepancy. The sponsor has  
21 now told us that they actually have improved upon that  
22 imbalance, and they're better when the way it was when

1 they had the -- during the study and now the FDA has  
2 reviewed this and said that there are some gaps  
3 between what the sponsor is proposing and what had  
4 actually gone on during the trial.

5           Let's focus again on those gaps and your  
6 interpretation of what the sponsor had just pointed  
7 out, that they've improved upon this. Have they  
8 actually improved this or do the gaps still remain  
9 between what was done in the trials and what's being  
10 proposed going forward for the use of this procedure?

11          And this is not just an injection. This is not  
12 somebody getting an injection into a de Quervain's  
13 tendon. This is a procedure where it involves  
14 manipulation following the injection and a recognition  
15 that a tendon might have ruptured or that a ligament  
16 was ruptured following the procedure, which requires  
17 some definite cerebral expertise in being able to sort  
18 that out postoperatively or postinjection.

19           So it's not just an injection. I think we  
20 need to keep that in mind as well.

21           DR. OKADA: This is Sarah Okada. So we did  
22 have an opportunity to review the revised training



1 manual and DVD that's proposed for use in clinical  
2 practice, and we actually concur with the sponsor that  
3 they've made some significant improvements in these  
4 things. And so they're fairly comprehensive.

5           The situation obviously is still somewhat  
6 questionable in terms of how much sort of hands-on and  
7 person-to-person training went on during the trials  
8 versus what would be the case during clinical  
9 practice. That's not so clear. Obviously, they're  
10 proposing to have some liaisons available. We're not  
11 completely sure what the background of those liaisons  
12 would be, whether they'd be available for any  
13 clinician who wanted to inject it and needed some  
14 hands-on assistance. Those details were not --  
15 haven't been finalized, so we're not clear on.

16           DR. RAPPAPORT: Ultimately, we're going to  
17 turn it back to you, Dr. Weisman, because it's really  
18 the questions that we're asking you today is based on  
19 the information we have, which is everything you've  
20 seen. We're not hiding anything. What do you think  
21 about whether the training is adequate, whether you  
22 think that the practitioners need to be from certain

1 groups or have a history?

2 All of those questions are what we're asking  
3 you, because there's no simple answer here and there's  
4 no way to study that without another ten years of  
5 extensive clinical trials that I'm not sure can even  
6 be done. In the meantime, we've got patients who may  
7 benefit from this.

8 DR. O'NEIL: Dr. Saag.

9 DR. SAAG: I think Dr. Rappaport and Okada  
10 are circling around the question that I asked earlier,  
11 and it may be that we're in a bit of somewhat  
12 uncharted territory with these risk management plans.  
13 But I think what would really help the panel out is to  
14 have a little more guidance from the FDA about what  
15 are the possibilities. Certainly, what the sponsors  
16 are proposing is reasonable. It's necessary, but is  
17 it sufficient? Is it enough?

18 We know from adult learning theory and other  
19 approaches to trying to train practitioners that you  
20 can increase knowledge but you may not change  
21 practice. You may not actually improve skills. So is  
22 this sufficient? It's necessary, but is it

1 sufficient? And understanding better what in this  
2 sort of new model that the FDA has adopted to more  
3 extensive risk management plans, knowing what other  
4 things are available in the armamentarium that the FDA  
5 could require would be very helpful to the panel.

6 DR. RAPPAPORT: I hear two questions in  
7 there. One is what can we do under our REMS, and the  
8 other maybe what is needed, or do we fully understand  
9 what's going to work in this situation and how we're  
10 going to assess that.

11 So with the second question, I'm not sure we  
12 have an answer to that yet, that we really don't know  
13 a lot yet about how REMS work. There's a lot of  
14 history of education, patient education, physician  
15 education. There are experts at the table here who  
16 can tell you more about it that we can probably.

17 But as to whether we should be imposing the  
18 restrictions that Kathy went over with you, and that's  
19 the limit of our restrictions. We can require that  
20 only certain prescribers, specialties, are actually  
21 doing these procedures or we can do nothing. Those  
22 are the extremes.

1           Let's go back to the fact that we have over  
2 the last couple of years since we've had this  
3 authority learned some new things about imposing  
4 restrictions, and you need to take the impact of  
5 imposing restrictions into consideration in whether we  
6 should really be doing that.

7           The company has already provided quite a  
8 restrictive plan, and whether it's going to work is  
9 yet to be seen. And, perhaps, what one possibility is  
10 to let them take the responsibility at this point for  
11 making sure that the right people are receiving or are  
12 being allowed to use the product. And then we can  
13 monitor over time and see how that's working. That's  
14 one option.

15          The other option is that we could step in  
16 and do our own mandated restrictions that have the  
17 authority of law and that we could fine people for not  
18 doing. If we do that, however, we're imposing a huge  
19 burden, and that's part of what was in that law, as  
20 Kathy explained to you, that we're not -- we have to  
21 consider how much of a burden we're placing on the  
22 healthcare system. There's a huge burden and a huge

1 price for every restriction that's put on any kind of  
2 medication.

3           If you think about what's out there, what's  
4 approved for use, there isn't any medication out there  
5 that doesn't have significant risks. Drugs are  
6 unsafe, and you have to consider what -- how far you  
7 want to go in having the federal government actually  
8 be the restricting agent. If this doesn't work, we  
9 can always step in later. If we see increasing  
10 problems with tendon rupture over time that are at a  
11 greater incidence than in surgeries or other new  
12 problems, we then still have the authority to step in  
13 and provide additional restrictions.

14           Did that answer your question at all?

15           DR. SAAG: It's helpful. Thank you.

16           DR. O'NEIL: Dr. Buckley.

17           DR. BUCKLEY: I guess two comments. One is  
18 that the company's provided a certain bar to get  
19 access to performing this in terms of your background  
20 training, but they really haven't talked about the  
21 volume issue, how many procedures do you need to  
22 perform a year, so that certain -- probably hand

1 surgeons are by definition doing that quite a lot, but  
2 rheumatologists might be performing one procedure a  
3 year or one procedure every two years. And if you  
4 believe it's not just your training but continued  
5 practice, and I think for most practitioners, it's  
6 probably continued practice where they learn 80  
7 percent of what they know. The training is 20  
8 percent. It's doing it over and over again is the  
9 other 80 percent. So we may be missing that in what  
10 they're offering us.

11           The other thing I have a little bit of  
12 concern about is, you weigh risk and benefit is a  
13 little bit of the benefit issue, because when we  
14 talked about that, we talked -- if you look at Slide  
15 10, where we talk about the success of getting to the  
16 primary endpoint by kind of training, one of the  
17 things that struck me was that although there wasn't a  
18 difference by the types of physician training, there  
19 was a big difference in success rate between Study 57  
20 and Study 59.

21           This was a placebo trial, but I imagine  
22 given that almost all the patients got adverse

1 reactions, neither the patient nor the physician was  
2 blinded for very long. I suspect that these  
3 measurements were made by the physician who did them,  
4 not an independent monitor or a picture taken.

5           So these results in the first study, 57, the  
6 good response rate was 60 to 70 percent. In 59, it  
7 was 40 to 47 percent. In the real world, maybe among  
8 people who are a little less experienced in this, the  
9 results may be less. So we have a procedure that no  
10 one denies probably is going to be very helpful to  
11 some people, and we have some risks that we don't  
12 quite understand. And when we put those things  
13 together, I think there's still some concern.

14           DR. DELCONTE: Dr. O'Neil, would I be able  
15 to address the issue of the differences between the  
16 studies, perhaps to shed some light on that?

17           DR. O'NEIL: If you can do it in under a  
18 minute. We're running a bit late already, and we have  
19 three more questions.

20           DR. DELCONTE: The difference can be  
21 explained by the difference in the severity, and I had  
22 shown earlier that the MP joints of low severity

1 performed better than the high severity. In Study  
2 857, most of the joints were MP of low severity. In  
3 the second study, in the Australian study, there was a  
4 predominance of PIP of greater severity, so that could  
5 account for some of the differences.

6 DR. O'NEIL: Thank you.

7 Dr. McAlindon.

8 DR. McALINDON: Thank you. So I'm just a  
9 little concerned about what I perceive to be a slight  
10 logic gap, in that we have great concern about the  
11 incidence of tendon rupture, and we're responding to  
12 it through restriction of access to individuals who  
13 have skill in hand surgery. But I think the numbers  
14 are too small in terms of that adverse event to really  
15 inform us one way or another whether any part of that  
16 risk was mediated by skill level. Indeed, the  
17 ruptures occurred among physicians who were presumably  
18 quite skilled at performing hand injections.

19 So it could be that the risk is mainly  
20 mediated by patient and disease characteristics rather  
21 than the skill level. In other words, it may be  
22 sufficient just to put the intervention in the right



1 place, and the rest of the consequences are then  
2 dictated by the patient characteristics.

3           So if that's the case and this is  
4 hypothetical, if that's the case, trying to mitigate  
5 that risk by a complex program that either restricts  
6 access or educates physicians might not in fact have  
7 much impact on the incidence of that outcome. And I  
8 think we just need to understand it better in order to  
9 figure out how to reduce that.

10           DR. SAAG: Can I respond to that?

11           DR. O'NEIL: Sure.

12           DR. SAAG: Tim, I agree with your premise,  
13 but I'm not sure that the clinical trials address the  
14 issue. I think there's a problem, that of  
15 generalizability; namely, all of the people performing  
16 the clinical trials were skilled. But what we don't  
17 know is what happens when we get out into the real  
18 world and we have people that spent 20 minutes  
19 watching a video doing this procedure who aren't  
20 familiar with the hand anatomy properly?

21           And knowing that there is the potential for  
22 a risk and that the risk has been seen in the clinical

1 trial, whether it's related to the patient  
2 characteristics or to the injection technique is not  
3 known, but one could easily speculate that if there is  
4 a component of injection technique that is in some way  
5 responsible for some proportion of adverse outcomes,  
6 that this would be manifest and magnified considerably  
7 greater in a real world setting than in a very  
8 controlled clinical trial.

9 DR. O'NEIL: Ms. Aronson.

10 MS. ARONSON: I have, I think, a quick point  
11 of clarification. I'm trying to understand what might  
12 fall under REMS. If there was some kind of guidance  
13 that if a patient presented as complicated; in other  
14 words, if the patient also had rheumatoid arthritis  
15 and some deformity, then it might be advisable for the  
16 patient to be referred to someone with high experience  
17 in hand issues. Would that fall under the REMS or,  
18 would that be just some guidance that could be put  
19 out?

20 DR. OKADA: This is Sarah Okada. I'm not  
21 really familiar with any REMS that gets down to that  
22 level of detail in terms of trying to dictate sort of

1 clinical practice. However, if it were really  
2 important and we thought that that would mitigate a  
3 specific risk, I could envision that some component of  
4 a REMS could be constructed to address that. I'm not  
5 certain that that's going to be the case here, though.

6 DR. RAPPAPORT: And it doesn't have to be as  
7 part of a REMS, either. It could be part of the  
8 program that the company is providing without us  
9 intervening.

10 DR. O'NEIL: We have time for one more  
11 question, and Dr. Haque has a question.

12 DR. HAQUE: I just have one question for  
13 Dr. Rappaport, and that's rather than go in the  
14 full-blown REMS with the cost and other issues that  
15 you mentioned earlier, can we -- are we in a position  
16 to make some suggestions that at least a registry be  
17 maintained so that we can quickly catch a trend if we  
18 see one rather than just have these patients get their  
19 doctors certified, they get their injection, they get  
20 one follow-up maybe, if that and then they're lost  
21 afterwards?

22 DR. RAPPAPORT: There's a whole range of

1 possibilities here, and we are very interested in what  
2 your thinking is on those, and we'll certainly take  
3 that into consideration in where we end up with this.

4 DR. O'NEIL: Well, I'd like to thank  
5 everyone for a lively and interesting discussion which  
6 we will be able to continue somewhat later in the  
7 program. We will now break for lunch, and we'll  
8 reconvene again in this room in 45 minutes, at 12:45.  
9 Please take any personal belongings you may want to  
10 with you at this time, and Committee members, please  
11 remember, again, that there should be no discussion of  
12 the meeting during the lunch among yourselves, with  
13 the press or with any member of the audience.

14 Thank you.

15 (Whereupon, at 11:57 a.m., a lunch recess  
16 was taken.)

17

18

19

20

21

22

2 DR. O'NEIL: Thank you, and welcome back to  
3 the meeting of the Arthritis Advisory Committee. We  
4 are about to begin the open public hearing, and before  
5 I do that, Nicole Vesely has some comments.

13           For this reason, the FDA encourages you, the  
14   open public hearing speaker, at the beginning of your  
15   written or oral statement to advise the Committee of  
16   any financial relationship that you may have with the  
17   sponsor, its product and, if known, its direct  
18   competitors. For example, this financial information  
19   may include the sponsor's payment of your travel,  
20   lodging or other expenses in connection with your  
21   attendance at the meeting.

1 beginning of your statement to advise the Committee if  
2 you do not have any such financial relationships.

3 If you choose not to address this issue of  
4 financial relationships at the beginning of your  
5 statement, it will not preclude you from speaking.

6 The FDA and this Committee place great  
7 importance in the open public hearing process. The  
8 insights and comments provided can help the agency and  
9 this Committee in their consideration of the issues  
10 before them. That said, in many instances and for  
11 many topics, there will be a variety of opinions. One  
12 of our goals today is for this open public hearing to  
13 be conducted in a fair and open way, where every  
14 participant is listened to carefully and treated with  
15 dignity, courtesy and respect. Therefore, please  
16 speak only when recognized by the Chair.

17 Thank you for your cooperation.

18 DR. O'NEIL: We will begin with comments  
19 from Mr. Tom Fewell. Mr. Fewell.

20 MR. FEWELL: Thank you for the opportunity  
21 to come here today to tell you about a treatment that  
22 changed and restored my life. My name is Tom Fewell.

1 I'm from Sycamore, Illinois. I was first diagnosed  
2 with Dupuytren's contracture in 1997, and have lived  
3 and adapted to the effects and limitations of this  
4 disease since then. Between my right and left hands,  
5 I've had two surgeries, and I've also participated in  
6 the AA4500 clinical trial in '07 and '08.

7 I received minimal compensation for mileage  
8 during the trial from Auxilium. I also participated  
9 in an advisory board meeting in November of '08, and  
10 Auxilium arranged and paid for the travel and lodging  
11 for that meeting. They also arranged for the travel  
12 and lodging so I could attend this meeting today. I  
13 do not own any stock in Auxilium.

14 Previous to my diagnosis in '97, I steadily  
15 lost function in my left hand. That affected my  
16 productivity at work, especially keyboarding and  
17 picking up objects. I also could not participate in  
18 recreational activities like playing ball with my kids  
19 or playing golf.

20 My first surgery was an outpatient  
21 procedure, and it was effective. But within months,  
22 more cords started to form on the ring finger of my

1 left hand, and I had surgery again on my left hand in  
2 2001. The second procedure was much more extensive  
3 and invasive to my nerves and skin. Anesthesia was  
4 used during the procedure, with its inherent risks. I  
5 lost time and productivity from my job during this  
6 time because of the follow-up appointments and six to  
7 eight weeks of rehab it took to regain my strength and  
8 flexibility.

9           So the surgery was more effective and  
10 long-lasting, but was also significantly more risky,  
11 painful and expensive.

12           As more cords started to form, this time in  
13 my right hand, I put off and delayed more surgery  
14 because of the pain, risk and expense until the  
15 contracture seriously interfered with my daily  
16 activities and ability to perform my job, as well as  
17 limited my personal life and mental outlook. Again, I  
18 lost productivity at work because of the difficulty  
19 and frequent mistakes in using a keyboard.

20           During my working career, I was a senior  
21 buyer, an international business buyer, and met  
22 business leaders through Asia, Europe as well as the



1 U.S. and Canada. Making a good impression is  
2 important, and that starts with a confident, firm  
3 handshake. As my hand contracted, I felt less  
4 confident and sometimes embarrassed because of my  
5 strange handshake.

6           Gardening was also difficult and painful.  
7 My hands tired easily when I played golf. Putting on  
8 gloves became a five- to 10-minute ordeal because I  
9 could not open my palm to insert my hand into the  
10 gloves, even mittens.

11           When I finally made the appointment to  
12 discuss surgery again in 2007, Dr. Beher (?) told me  
13 about an upcoming clinical trial using injections and  
14 outpatient type of procedures rather than surgery. I  
15 was in a unique position to be able to compare and  
16 contrast the effectivity and risk of injections versus  
17 surgery. Two surgeries, one set of injections, it was  
18 an easy choice.

19           I participated in the trial. During the  
20 treatments, I did feel very sharp pains during the  
21 10-second cycles when the doctor physically  
22 straightened my hand, breaking the cord tissues apart.

1 I managed coping with those intense cycles because I  
2 knew and expected that when the cord tore apart, I  
3 would experience immediate and significant improvement  
4 in my hand movement, range and flexibility. And after  
5 three cycles, I recovered full use of my hand.

6 I engage in all activities I want to,  
7 recreational, keyboarding, the many things you take  
8 for granted, and putting your hands in your pockets to  
9 get keys, I could do that again. And I could shake  
10 hands, and I could clap when my kids did something  
11 well, something I couldn't do before.

12 The collagenase treatments eliminate the  
13 risk and uncertain results of surgery. The treatments  
14 resulted in immediate and effective improvement in  
15 hand motion and quality of life. There is some pain,  
16 but it is manageable, especially considering the  
17 extensive rehab that is also eliminated. And although  
18 longevity is still being studied, my treatments ended  
19 18 months ago and the cord has not reappeared. I  
20 think the injection treatment represent a win-win  
21 scenario for reducing pain and improving the quality  
22 of life and reducing patient risk.

1                   Thank you.

2                   DR. O'NEIL: Thank you, Mr. Fewell.

3                   The next speaker will be Rodney Van Sickle.

4   Mr. Van Sickle.

5                   MR. VAN SICKLE: Hi, my name is Rod Van

6   Sickle, and I've had Dupuytren's for about 12 years.

7   I do not own stock in Auxilium, and I've received only  
8   minimal compensation for my time and travel.

9                   Dupuytren's affected my right hand first. I  
10   tried to postpone it because the doctor told me at  
11   that time that there was a drug going to become  
12   available, but I couldn't wait any longer. I couldn't  
13   perform my job as a fire captain any longer, so I went  
14   ahead and had the surgery.

15                  After the first surgery, I got a staph  
16   infection, and that required me to have a second  
17   surgery. And the disease came back with a vengeance  
18   in my little finger in my right hand, completely  
19   closed against the palm of my hand, and I had to have  
20   a third surgery. The little finger on my right hand  
21   now is at about 90. This is as good as it gets.

22                  When the trial became available and I was

1 allowed to be in the trial, and after the three  
2 injections -- before that, the ring finger on my left  
3 hand was at about 50 degrees. After the three  
4 injections into the cord on that hand, the cord popped  
5 after the third injection, and my left hand is  
6 perfectly straight.

7           And if you compare the -- there is no  
8 comparison between -- in my opinion, between the  
9 surgery and the drug injections. It's just -- I had  
10 such a horrible time with the surgery, nothing against  
11 the surgeons, but it just didn't work well for me.

12           The disease runs in my family. My father  
13 has it. My brother has it, and my two sons have it to  
14 different degrees. And I would really like to urge  
15 you guys to approve this drug for the market because  
16 it would help so many people. Thank you.

17           DR. O'NEIL: Thank you.

18           The next speaker is Ms. Karen Mercaldo.

19           MS. MERCALDO: My name is Karen Mercaldo.  
20 I'm 61 years old. I do not own stock in Auxilium, and  
21 I've not been compensated except for my travel  
22 expenses.

1           I was diagnosed with Dupuytren's disease in  
2 1996 during surgery for what the doctors thought was a  
3 cyst. I was 47 years old at the time, and I knew what  
4 Dupuytren's was because my father has had it ever  
5 since I can remember. He had two unsuccessful  
6 surgeries in the '50s, and as a child, I remember  
7 hearing him say that he would cut his hand off before  
8 he ever went through that again.

9           My recovery from surgery was long and  
10 difficult. For the first week, I had to wear a sling  
11 that positioned my hand directly in front of my face.  
12 For months, I kept dropping things and sometimes  
13 burning myself because it took that long to get the  
14 feeling back in my hand.

15          Within two years, the problem returned on  
16 the same finger and two others. It was much worse  
17 than before, and on both hands. I was discouraged.  
18 It didn't seem worth going through the surgery if the  
19 condition was going to return. Three of my fingers  
20 were affected at the PIP joint so that my hands looked  
21 like this.

22          Everything I did became more difficult.

1 Simple things like putting on a glove or typing were  
2 cumbersome. I managed to adjust in many ways, but had  
3 to give up completely things that had enriched my  
4 life, such as playing the piano, knitting and  
5 painting. I couldn't even wear my wedding ring.

6           Then one night a friend of mine who also has  
7 Dupuytren's showed me his pinky, which was almost  
8 completely straight. He said to me, "Yesterday, it  
9 was like this."

10           He told me that he was a patient in the  
11 clinical trial, and I didn't need any convincing to  
12 find my way there. I had my injections in August,  
13 October and November of 2003. The morning after the  
14 first injection, the doctor took my finger and  
15 straightened it. It hurt momentarily, but the tears  
16 in my eyes were tears of joy. I was thrilled. I  
17 found that although there was some soreness, I could  
18 use my hand immediately. I did some exercises and  
19 wore a splint at night. I anxiously awaited the  
20 subsequent injections and was just as pleased with the  
21 results.

22           Now, six years later, I still have the full

1 use of both of my hands. I enjoy playing the piano.  
2 I've started a knitting club for my granddaughter and  
3 her friends. I never poke myself in the eye while  
4 washing my face anymore, and I'm wearing my wedding  
5 ring. I even started playing the viola again after  
6 many years, something I thought I would have the  
7 dexterity to do. I enjoy it immensely, and I play in  
8 church every week.

9           My Dupuytren's symptoms have not returned,  
10 but more significant to me is that my fear of the  
11 symptoms returning, which was very great, is gone.  
12 Most people never think about the blessing it is to  
13 have the use of their hands, but I think about it  
14 every single day, and every day, I'm thankful. I have  
15 a son who is a gifted pianist, and grandchildren who  
16 show promise in music and art. It is for my children  
17 and grandchildren and not just for myself that I  
18 appeal to you to approve the drug Xiaflex. It would  
19 mean so much to me, to my family and to the many  
20 others who suffer from this debilitating condition.

21           Thank you.

22           DR. O'NEIL: Thank you, Ms. Mercado.

1           The next speaker is Kenneth Nelson.

2           MR. NELSON:   Good afternoon.   Madam  
3   Chairman, thank you, and the Committee, for your time  
4   today and for the opportunity to share my experience  
5   as both a victim of Dupuytren's contracture and as a  
6   beneficiary of medical research that is the subject of  
7   today's hearing.   My testimony is mine and mine alone.  
8   I do not own any stock in Auxilium Pharmaceuticals.   I  
9   have not been paid anything for my appearance today  
10   other than for travel, lodging and meals, and a small  
11   stipend for my participation in the clinical trials.

12           I have no other motive for being here today  
13   other than to voluntarily share how this exciting has  
14   returned a quality of life I really enjoyed but I lost  
15   more than 20 years ago.   I first became aware that  
16   something was wrong in my early 30s.   I noticed some  
17   lumps developing in a pit in the palm of my left hand.  
18   By my early 40s, these things appeared in both hands.  
19   It felt like a cord or lumps just under the skin.  
20   There was no pain.   I had no restrictions of movement  
21   in my fingers or thumbs, but it was something that I  
22   brought to the attention of my family physician.



1 Well, since I did a lot of physical work around the  
2 house, he passed those lumps off as calluses and we  
3 left it at that.

4 But a few years later, I noticed that my  
5 third and fourth, or ring and small fingers and  
6 thumbs, were slowly beginning to contract. By age 45,  
7 the condition was beginning to prevent me from doing  
8 things that I loved to do like playing the piano, or  
9 as it's even been said here, washing my hair without  
10 poking myself in the eye, or even trying to put on a  
11 pair of gloves or shaking hands with clients.

12 My doctor then finally referred me to the  
13 Indiana Hand Center. The diagnosis was Dupuytren's  
14 contracture. I had not a clue what that meant. I was  
15 told the condition would only worsen, that the only  
16 option at that time was radical surgery. I was  
17 shocked to see the extent to which my hands had to be  
18 cut or sliced open to remove the growth, followed by a  
19 boxing glove-type wrap and lengthy rehabilitation to  
20 regain movement and strength. There was also the  
21 possibility that I could lose some of the feelings in  
22 my fingers. There was the threat of potential

1 infection and that Dupuytren's was likely to return.

2           Radical surgery was not an option that I was  
3 willing to accept at that time. At least I could  
4 still hunt and peck on the computer keyboard. I even  
5 found sort of a simple way to play Chopsticks on the  
6 piano again. But this continued to worsen and  
7 seriously affected my quality of life. Even when  
8 shaking hands with people, they would often ask me,  
9 Ken, have you been the victim of a stroke -- were  
10 embarrassed to ask you. I would jokingly tell them as  
11 I held their hand that I had a highly contagious  
12 disease and they would withdraw very quickly, but I  
13 tried to find some humor in these conditions.

14           I remained hopeful that medical science  
15 would develop a safe, noninvasive procedure that would  
16 give me back the use of my hands without having to go  
17 under the knife. Well, that day came when my wife  
18 Susie read a notice in the newspaper seeking  
19 candidates for a clinical trial designed to reverse  
20 the devastating impact of Dupuytren's. After  
21 undergoing a medical exam and extensive questioning, I  
22 qualified as a candidate.

1           Dr. Kaplan, who is here, of the Indiana Hand  
2 Center, was very thorough in explaining the process  
3 followed by the cord rupture procedure. He made sure  
4 I understood the process, beginning with the need for  
5 blood chemistry. My first injection was uncomfortable  
6 but tolerable. Attempts to rupture the cord proved  
7 futile and painless. I was among the group to get the  
8 placebo.

9           Well, eventually, I received the real thing.  
10 Once again, the injection was uncomfortable, this time  
11 with an increased stinging sensation. My fingers were  
12 immobilized with a wrap, and I was sent home. Later  
13 that day, I accidentally bumped my fingers. I felt  
14 something like a wasp or bee sting in the palm of my  
15 hand. I already could begin to hear this cord  
16 rupturing or popping. I could feel it. Well, upon  
17 going to bed that night, I removed the wrap as  
18 instructed and noticed a slight bruising in the area  
19 of the injection. I was very careful not to extend my  
20 fingers or intentionally try and rupture the cord,  
21 although it was tempting.

22           The next morning, I returned to the hand

1 center. Because of the trial requirements, numbing  
2 medication was not administered, and Dr. Kaplan  
3 explained there would be a rather sharp pain when and  
4 if the cord ruptured. But it would last only a  
5 moment. He was right. As Karen has said, when I heard  
6 that cord pop and saw my fingers suddenly straighten  
7 out after years of being jammed into the palm of my  
8 hand, tears came to my eyes. I get emotional about  
9 this still.

10           It was partly because of the moment of pain  
11 but mostly due to the emotion of witnessing what I  
12 still call a miracle in my life. Well, at the end of  
13 each rupture session, I was fitted with a night brace  
14 to help keep my fingers from retreating back into the  
15 palm of my hand.

16           I can just simply sum up my remarks by  
17 saying that I inherited this from my dad who has  
18 Dupuytren's. Our youngest son Bradley is beginning to  
19 get the pits and the cords in the palm of his hand.  
20 It's worked for me. It's been a wonderful procedure.  
21 I thank           Dr. Kaplan for his professionalism.  
22 And I highly encourage you to go ahead and proceed

1 with this and give it an okay.

2 Thank you.

3 DR. O'NEIL: Thank you.

4 The next speaker is Bill Walker.

5 Mr. Walker.

6 MR. WALKER: Hello, ladies and gentlemen. I  
7 want to start with the disclaimer that I have no  
8 financial affiliation with Auxilium, and I've just  
9 been also paid -- not paid but just reimbursed for  
10 travel expenses. I'm here today purely out of my  
11 enthusiasm for this drug because it changed my life  
12 totally. I get emotional, too. I'm sorry.

13 But like Ken, it started in the 30s. You  
14 get a pitting in your hand, and you don't know what it  
15 is. And over time, it's very insidious, and it takes  
16 maybe eight to 10 years to really where it draws your  
17 hand back to -- it's not useless, but it's close to  
18 that. I mean, you can drive and you can live, but you  
19 can't live like you used to. You can't play tennis.  
20 You can't -- again with gloves, I'm a device rep, and  
21 I work in the OR in a lot of hospitals. And you can't  
22 even put latex gloves on to protect your hands

1 from -- you know universal precautions. You would  
2 have to get a sterile towel and grab a cable that is  
3 passed off to you because it has come from -- it's a  
4 surgery thing.

5 But anyway, being included in the trial, I  
6 was on vacation two years ago. We were in Italy, in  
7 Tuscany, and my tour guide was from Indianapolis. She  
8 was in Dr. Kaplan's trial. She looked at my hand and  
9 said, "You have Dupuytren's. You should go to the  
10 hand center and see if you can be in this study."

11 So I met Dr. Kaplan, and he looked at it. I  
12 was fortunate enough to be randomized to the real  
13 deal. We had the collagenase enzyme. The first  
14 injection, that night at home watching TV, my hand  
15 starts to pop open just spontaneously with the boxing  
16 glove dressing on, and it was nothing short of  
17 miraculous. And I go back in and see Dr. Kaplan the  
18 next day. My hand was very affected. It was 70  
19 degrees back to -- these two fingers on the right  
20 hand. Within the -- after the first month, you go  
21 back and you get another round of injection. It was  
22 virtually straight at that time, one month and the

1 hand's almost normal.

2 I had seen one of his colleagues maybe a  
3 year prior to that, and like Ken was saying, he saw  
4 the surgery and it's a lot. It's very scary to have  
5 to think about the surgery. I work probably 70 hours  
6 a week, and I don't have time for surgery. I don't  
7 have time for physical rehab because I'm very devoted  
8 to my job.

9 But anyway, being included in this study has  
10 really changed my life. And if you also look at if  
11 you go to surgery, the anesthesia risk, too. That's a  
12 big risk, the risk of infection, which you heard Rod  
13 say he had that happen, too. And with the enzyme  
14 injection, they target -- Dr. Kaplan was really --  
15 well, he's an expert. He can look at the cord, a very  
16 fine needle penetrates into the tissue, three  
17 injections, 0.58 milligrams and it -- being an enzyme,  
18 it just dissolves the tissue and allows the hand to  
19 break open.

20 Now, if you compare that to a surgical  
21 approach, it's just -- the surgical approach is a  
22 totally different animal, and the benefit from the

1 injection I think just outweighs by far a surgical  
2 approach.

3           And the things about -- I drive sometimes  
4 240 miles a day. I would get -- I would see him in  
5 the office at 7:00 a.m. I could be in the OR by  
6 10:00 in the morning at Madison, Indiana and be  
7 performing my work. Even with a dressing on my hand,  
8 you can still work around that, and the dressing is  
9 only on for 24 hours. But I lost no time at work, and  
10 I had a perfect result.

11           This hand was 70-plus degrees back. It's  
12 straight as an arrow now. It's been that way for a  
13 year. And at Christmastime, you get gifts like  
14 gloves, my mother, she'll get you gloves, and I could  
15 never wear them. When she saw my hand, she had tears  
16 of joy. She was so happy. And it's just -- I'm really  
17 privileged to be part of the program here. And I just  
18 want to thank the hand center and Dr. Kaplan.

19           And you folks in the FDA, we need this out  
20 in the streets as soon as we can, so I'm a total  
21 believer in it and a recipient as well. And I just  
22 want to thank you all.



1 DR. O'NEIL: Thank you, Mr. Walker.

2 We will now have very brief comments from  
3 Dr. Robert Hamilton, a Ph.D. immunologist from Johns  
4 Hopkins.

5 DR. HAMILTON: Thank you, Panel.

6 First, I own no stock in Auxilium, and I  
7 have no vested interest in the drug itself one way or  
8 the other. I'm here today because my clinical  
9 laboratory at Johns Hopkins did the initial  
10 immunogenicity studies on the Phase 1, Phase 2 and  
11 early Phase 3 studies of Dupuytren's that were done at  
12 Stony Brook back from 2001 to 2006.

13 And as you who are medically qualified know,  
14 there are five classes of immunoglobulin or  
15 antibodies, and of those, IGE drives allergic disease  
16 and IGG is viewed more as protective. So one of my  
17 puzzles was not to see the breakdown of the immune  
18 responses in the Phase 3 study into IGE and IGG.

19 So in our initial testing with the  
20 Dupuytren's sera from Stony Brook, what I can say is  
21 that we detected IGE antibody to collagenase in  
22 approximately a third of the individuals who were

1 subjected to the analysis or to the studies. After  
2 repetitive injections, some of these levels arrived at  
3 levels that we see with patients who have hymenoptera  
4 venom allergies and have reactions.

5           Because this was a primary immune response,  
6 you would not expect to see allergic reactions during  
7 the first three months of treatment, because the  
8 immune response is just beginning. The concentration  
9 is low. The affinity is low. The specificity is not  
10 what it could be.

11           And the Phase 3 study clinical data today  
12 support the notion that in fact, the first course of  
13 treatment of three injections is safe. It doesn't  
14 elicit obvious systemic reactions. So up to one to  
15 three injections -- based on the data, the clinical  
16 data supports the safety of it.

17           Today, we heard that there was 100 percent  
18 of Dupuytren's patients who elicited antibody  
19 responses. I'd like to know how many of those elicited  
20 IGE, not because in the first course of treatment they  
21 would be expected reactions, but if they ever choose  
22 to come back for a second course of treatment, that's

1   precisely where you're going to see the systemic  
2   reactions.  And I would suggest that if you do license  
3   the drug, that you license it for a first course of  
4   treatment, and that you request additional studies to  
5   document the safety of the drug when patients come  
6   back for repetitive administrations four to six months  
7   after administration.

8               Second, that you identify or define what is  
9   a large local or a systemic reaction so they know what  
10   to look for, and that any individual who manifests  
11   those symptoms in fact gets a blood sample and gets at  
12   least evaluated from an immunogenicity point of view  
13   in terms of IGE and IGG antibody responses that are  
14   technically capable -- we're capable of doing those  
15   measurements analytically today.

16              So thank you very much.  I only have three  
17   minutes.  Thanks.

18              DR. O'NEIL:  Thank you.

19              We had a number of questions left over from  
20   the first session, questions to the sponsors from the  
21   panel members, and we will begin there with Dr.  
22   Kaplan.

1 DR. S. KAPLAN: Thank you. As a hand  
2 surgeon who's dealt with this condition for many  
3 years, I welcome a viable alternative to surgery, and  
4 this may be such an alternative.

5 I do have concerns. I share everybody's  
6 concern about the crucial nature of the injection, and  
7 making sure that the right people who know the  
8 condition and understand are involved.

9 I share Dr. Swartz's concern about off-label  
10 use. I know that there are Stage 2 clinical trials  
11 underway for use in frozen shoulder and Peyronie's  
12 disease. At a recent medical meeting, if you stopped  
13 by a booth, you got a candy bar and you can do a  
14 survey, and the survey was clearly about this use of  
15 this product in other conditions.

16 Severe scarring being one of them, plantar  
17 fasciitis being another. I can envision a variety of  
18 conditions where people might want to try this  
19 off-label, and I would be worried about that.

20 I have two very specific questions. One,  
21 you mentioned an ongoing study of two- to five-year  
22 follow-up. I'm sure you are aware that the results

1 from Stanford of an eight-year follow-up recently  
2 presented at the American Society for Surgery, the  
3 hand meeting at San Francisco. They had eight people  
4 followed after eight years. Six of the eight had  
5 recurrence. In two situations, the recurrence was  
6 actually worse than on original presentation. The  
7 four others, it was mild and two others, it was -- it  
8 did not recur.

9           And although your two- to five-year studies  
10 are not complete, do you have other data on recurrence  
11 that you've not shared with us?

12           My second specific question involves your  
13 recommendation for injecting only one cord at a time.  
14 You demonstrated product safety. We just heard about  
15 some concerns about IGG and IGE. If the  
16 recommendation is to inject one cord at a time and you  
17 have what in my office is a fairly common situation of  
18 bilateral involvement with multiple fingers, it could  
19 conceivably be that a person would come over the --  
20 for 24 visits with 12 injections and take a year to do  
21 so for treatment of two fingers on each hand.

22           And my question is then with the safety

1 profile you outlined, are you going to modify that  
2 recommendation? Thank you.

3 DR. DELCONTE: Let me address the first  
4 question first about the recurrence, and then I'll  
5 have Dr. Tursi talk about the injection regimen,  
6 because there were a number of patients who had  
7 various intervals between injections.

8 What we had talked about was in the  
9 durability of response was 830 successful patients  
10 treated, we had 30 recurrences. And as you'd heard,  
11 the definition of that is a recurrence to a  
12 contracture of greater than or equal to 20 degrees  
13 with a palpable cord. If you do Kaplan-Meier  
14 estimates, the rates at one year are 6.7 percent on  
15 the successfully treated joints.

16 The data we showed from some of the surgical  
17 therapies -- and this is the average follow-up of 12  
18 months -- within or lower to that range, about 19 to  
19 22 percent.

20 Then the last question about the follow-up  
21 study, that's an ongoing study which will take the  
22 patients from the current clinical trials, and that's

1 a two- to five-year follow-up. So we'll get some  
2 additional long-term recurrence data there.

3 And that's -- to answer the question about  
4 the series of eight patients from San Francisco, we  
5 realize that's a small number of patients, and that's  
6 the reason why we'd like to do the long-term study.

7 And then, Jim, you want to come up and  
8 address?

9 DR. TURSI: Sure. Jim Tursi, with Clinical  
10 Affairs.

11 As to the safety of injecting more than one  
12 joint at a time, that would not be a recommendation  
13 that we have, as it was not studied during the  
14 clinical program. If you'd like to see, I can show  
15 you some details on subjects that were treated close  
16 together in proximity, meaning short inter-injection  
17 intervals. But I'll leave that at your discretion, if  
18 you'd like to see that.

19 DR. O'NEIL: If it's informative, I think we  
20 would like to see that.

21 DR. TURSI: Okay. What I've done is I've  
22 taken those subjects that have had essentially two

1 weeks or less between injections. And as you can see,  
2 these are the subject number along the left side. And  
3 the days between the injections ranges from 10 upwards  
4 to 15. Just to explain the organization of the table,  
5 at the top is the original joint that was treated, so  
6 in this case, it was the left ring PIP joint. And in  
7 parentheses, it just demonstrates that it was a  
8 success.

9           The joint below was the one that was  
10 ultimately treated at the interval following. So this  
11 was treated 13 days later. This particular joint 10  
12 days later.

13           And what's important to point out was that  
14 even in these subjects, first of all, they were all  
15 successful with these short intervals. The second  
16 important point is to point out that when you consider  
17 the adverse event profile, the adverse events that we  
18 saw in these subjects was no different than those  
19 subjects who had received at a 30-day interval or a  
20 longer interval.

21           DR. O'NEIL: Thank you.

22           We have at least two other questions for the



1 sponsor that have been identified to me, but I had  
2 promised the sponsor three minutes or so to present  
3 additional data. Are you ready to do that?

4 DR. DELCONTE: I just had one point of  
5 clarification on the qualifications and actually the  
6 training of our investigators. As we had mentioned  
7 before, we selected investigators who were hand  
8 surgeons, orthopedic and rheumatologists. And their  
9 relative level of experience, we had one  
10 sub-investigator who was in their first year out of  
11 fellowship, and we had several who had been in  
12 practice for more than 20 years.

13 Regarding the sites, we had not -- in  
14 addition to academic medical centers, we had large  
15 research clinics as well as private practices. So we  
16 tried to get as broad a range of possibility in the  
17 sub-investigators with regard to training and type of  
18 practice. That's all.

19 DR. O'NEIL: Thank you.

20 The next question was from Dr. McAlindon.

21 DR. McALINDON: It was a quick question in  
22 relation to, again, the tendon rupture issue. So I've

1    been wondering about whether tendon rupture that  
2    occurs as a consequence of AA4500 might be more  
3    difficult to repair than tendon ruptures that occur in  
4    other situations. I'm wondering if we have any data  
5    on that, or whether the operative findings perhaps  
6    were informative in that respect.

7               DR. DELCONTE: Yes, I'd like to ask  
8    Dr. Kaplan to come up and talk about operative  
9    findings on patients who've had AA4500.

10              DR. T. KAPLAN: There's a number of details  
11    in the three patients who had tendon rupture  
12    intraoperatively. I unfortunately had the opportunity  
13    to see one of them firsthand, as one of the tendon  
14    rupture patients was one of my own. That patient,  
15    unfortunately, he was also someone who had previously  
16    had surgery on his other hand. After that surgery, he  
17    experienced a flare reaction, and he was out of work  
18    for six months. So he was very interested in the  
19    potential for less-invasive treatment.

20              Unfortunately, with his first injection of  
21    Xiaflex given for the PIP joint, it was actually given  
22    at the radial base of his small finger. He went back

1 to work with limited time off and was moving a pallet.  
2 So we didn't -- at that point, he was the first point  
3 tendon rupture that had happened. There was no  
4 recommendation at that point that I had given him what  
5 not to do as far as forcible use of his hand.

6 He had a heavy pallet to move, was lifting  
7 up that heavy pallet with a pallet jack when he felt  
8 that tear.

9 Because of his experience with surgery on  
10 his other hand, he was very reticent to undergo a  
11 surgical procedure on that hand. So because he had  
12 ruptured his FDP tendon but his FDS tendon was intact,  
13 we first watched him to see whether or not he would  
14 function well with a superficialis finger, meaning  
15 that we didn't expect him to get motion back at his  
16 DIP joint, but he could still have functional motion  
17 at his PIP joint.

18 Unfortunately, he didn't get back the motion  
19 that he wanted. He had some discomfort from where the  
20 tendon rupture was. And when we explored him, found  
21 what we would typically see with probably more like a  
22 rheumatoid tendon rupture, where there was an

1 attritional rupture of that tendon. There were not  
2 healthy tendon ends to consider repair to, and the  
3 decision was at that point -- I talked to him  
4 beforehand about tendon grafting procedures versus  
5 just excision of the FDP remnant and a tenolysis of  
6 the FDS, which is what we did in his circumstance.

7           In one of the other tendon ruptures at one  
8 of the other sites, they intervened much more quickly,  
9 but again, they had to do a tendon reconstruction  
10 procedure, excise the damaged tendon and put in a  
11 tendon spacer for several months and then went back to  
12 do a tendon grafting procedure afterwards.

13           So I anticipate that when ruptures happen  
14 due to collagenase, that it would be a rupture that  
15 would not be directly repairable. You'd have to  
16 consider reconstructive options.

17           DR. O'NEIL: Thank you.

18           Dr. Olsen had a question as well.

19           DR. OLSEN: I had a question actually that  
20 was just touched on in the discussion, which was I  
21 wondered what the antibody classes were of the  
22 antibodies that the patients made to the drug, not

1 just IGE, but I was concerned also about classes of  
2 IGE that might consume complement and with re-  
3 challenge, you might face problems, for example, with  
4 immune complex formation.

5 DR. DELCONTE: Okay. I'll have Paul  
6 Chamberlain address that, and I should also mention we  
7 did have a number of patients, because of the way the  
8 trials were designed, who had a large interval between  
9 their double-blind portion and when they went into the  
10 open-label portion. So there was in some instances  
11 more than six months. And some patients in earlier  
12 trials had been exposed up to five years earlier,  
13 who'd been in later trials without any untoward  
14 effects.

15 So, Paul.

16 DR. CHAMBERLAIN: It's Paul Chamberlain, NDA  
17 Regulatory Science.

18 Yes, just to address the assay specifics.  
19 We were measuring total antibody. That's AUX-I or  
20 AUX-II specific antibody, regardless of class. So  
21 that would be substantially IGG, and we probably  
22 wouldn't detect specific IGE in the assay, but it

1 would be measured in the total assay.

2 I think the issue in terms of the IGE  
3 question that Dr. Hamilton asked is best addressed in  
4 terms of patients who followed up into Study 858 from  
5 857; that is, they had a series of treatments in one  
6 study and then rolled into a second study. And that  
7 would be when you most expect to see an exacerbation  
8 of the immune-mediated adverse drug events.

9 And these data show in the top panel the  
10 first pivotal study, this was 857, subjects on  
11 successive injection showed increasing titers of  
12 anti-AUX-I and anti-AUX-II antibodies as you move  
13 across from the first to the fifth injection.  
14 Subjects then rolled over into a follow-up study, an  
15 Open-label Study 858, and you can see at the time of  
16 the first injection, the titers were pretty much back  
17 down to the baseline level, but then rebounded on the  
18 second, third, fourth and fifth injections.

19 So this is exactly the scenario that  
20 Dr. Hamilton would expect to see, an exacerbation of  
21 immune-mediated adverse drug events. So I would like  
22 to hand over to my colleague Dr. Jim Tursi, just to

1 talk about the adverse events.

2 DR. TURSI: Recognizing that the antibody  
3 titers were higher in that specific study consistent  
4 with kind of the scenario that was described, I can  
5 take you through the adverse event profile  
6 demonstrating no difference, if not actually an  
7 improvement in the adverse profile of AA4500 in those  
8 subjects.

9 I'll take you through the same adverse  
10 events greater than or equal to 5 percent, and the  
11 left columns represent those in the 857 trial, the  
12 first trial, and the lighter green on the right side  
13 represent those subjects in the 858 trial with higher  
14 antibody titers. And what you can see across the  
15 adverse event profile is whether we consider swelling  
16 of the hand, contusion or injection site pain,  
17 extremity pain, hemorrhage, tenderness, et cetera, all  
18 those adverse events, right down to injection site  
19 pruritus -- there are no differences, if not  
20 improvements, in the adverse event profile looking at  
21 that trial of 857 to the trial with the higher  
22 antibody titers in 858, suggesting that there does not

1 appear to be any risk consistent with duration of  
2 injection.

3 I could also speak to subjects who have long  
4 interjection intervals, specifically those in the  
5 five- to six-year range, and what that information  
6 demonstrates is that there was no difference in terms  
7 of the adverse event profile in subjects even if they  
8 received it as far as ten years between injections.  
9 So, again, supporting the safety profile of AA4500 in  
10 the presence or absence of antibodies.

11 DR. O'NEIL: Could I quickly follow up and  
12 ask how many of the people in 857 did not roll over  
13 into 858, or was it a complete rollover? By that I  
14 mean people who did not go on to the follow-up study  
15 may have indeed been those who were at higher risk for  
16 some reason.

17 DR. DELCONTE: Only six of those patients in  
18 that study did not roll over out of the 308.

19 DR. O'NEIL: Okay. Thank you.

20 The one other question that I had is after  
21 there were three people who had tendon ruptures, you  
22 indicated that you changed the injection technique,



1 particularly for PIP joint injection or injection near  
2 the PIP joint for PIP contracture. Do we have any  
3 evidence whether that changed the outcome? That is,  
4 did the complication rate decline?

5 DR. DELCONTE: We've actually looked at the  
6 number of injections before and after. Dr. Tursi will  
7 go through that.

8 DR. TURSI: What we noticed with the  
9 training reinforcement was essentially that there was  
10 an improvement in terms of the potential risk that  
11 ultimately patients would foresee with the injection.  
12 This was a training reinforcement timeline, and  
13 essentially, ahead of the training reinforcement, we  
14 had performed 734 injections, 446 MP and 288 PIP  
15 cords. And as you can see, the two tendon ruptures  
16 occurred.

17 At the time of the training reinforcement,  
18 that was followed by over 1800 injections, 1,027 MP  
19 cords and 869 PIP cords, and a single tendon rupture.  
20 And I think it's important to point out that our  
21 injection training and the entire risk management  
22 program is designed with this experience in mind,

1 taking the lessons learned from the clinical trials  
2 and ultimately improving them for inclusion in the  
3 clinical program training.

4 DR. O'NEIL: Dr. Haque? I'm sorry, yes.

5 DR. HAQUE: Thank you. I'd like to wrap up  
6 a few of my last questions. First, you had about 13  
7 percent non-responders by the 50 percent improvement  
8 in contracture criteria, and 35 percent by the 5  
9 degree criteria.

10 Any thoughts on those non-responders, or are  
11 there any clues as to who we should not bother to  
12 inject?

13 The second question would be, is there any  
14 data on the safety of efficacy of surgery after the  
15 injection? Did any of your patients go on to need  
16 surgery, and was there any increased tissue damage  
17 present or any other problems with wound healing?

18 And then I had a question on your -- in the  
19 brief that we got before this meeting, I read the  
20 instructions that you were giving out, and it  
21 suggested that for the small finger, you would inject  
22 more towards the palmar digital crease. And I was

1 concerned about that, because that's where the spiral  
2 cords and abductor digiti minimi cords tend to push  
3 the neurovascular structures.

4 DR. DELCONTE: Let me start with dealing  
5 with what happened and what we saw in some of the  
6 non-responders. What we saw sometimes of the patients  
7 who didn't get down to zero to five, that they  
8 did -- there were a number that had some improvement.  
9 Of the ones that did not get all the way down to zero  
10 to 5 after -- or didn't get three injections, some of  
11 them did not have any more palpable cord, and the  
12 AA4500 was able to disrupt the cord, but it  
13 didn't -- there were other factors which may involve  
14 the collateral ligament, volar plate that could impact  
15 the finger from being completely straight.

16 Dr. Kaplan can talk about there were  
17 patients who were operated on after AA4500.

18 DR. T. KAPLAN: Yes, I guess I got to have  
19 experience with everything. For a while, I had one  
20 non-responder who had a really thick -- and I think  
21 that he didn't -- I ultimately took him to surgery  
22 because he didn't respond, and he just had a really

1 big, thick cord. And actually, when we got to  
2 surgery, you could see an area where that cord looked  
3 a little bit thinned superficially, almost like there  
4 was a little divot there. But the cord didn't break.  
5 So he just had a really thick cord. He -- we tried  
6 good, hard manipulations all three times, and he just  
7 didn't rupture.

8 I did have another patient who I took to  
9 surgery. She actually had eight injections. She had  
10 three placebo injections, followed by five collagenase  
11 injections. She had three collagenase to the MP joint  
12 level, two injections to the PIP joint level. And you  
13 can kind of see I have a free elevator, a little  
14 surgical instrument here, pointing to an area of the  
15 cord just to orient the fingers pointing out towards  
16 the left here. And the cord kind of comes up, and you  
17 can see the section from about here to about here no  
18 longer looks as well-defined as it does here, or even  
19 out here, although I think this area out here may have  
20 been the site of one of the PIP joint injections.

21 But this is the site where I think I did  
22 most of the injections, and you can see it just kind

1 of looks a little bit chewed up a little bit, a little  
2 bit reddened. It doesn't have that same organized  
3 consistency.

4 But speaking to the technical abilities to  
5 do that surgical procedure, I didn't find that the  
6 tissue planes were obliterated. It was still  
7 relatively easy to identify the fat layer from the  
8 neurovascular bundles, to safely identify the core  
9 tissue and excise it.

10 And I think there was a third clinical part  
11 that I forgot.

12 DR. HAQUE: The palmar digital crease  
13 injection for the small finger.

14 DR. T. KAPLAN: Yes, actually, I did a  
15 spiral cord.

16 DR. O'NEIL: Could you repeat the question?

17 DR. T. KAPLAN: Oh, I'm sorry. The question  
18 was at the base of the digit, as we modified the  
19 technique, and if I can just switch gears for a quick  
20 second, I think with the modified technique -- again,  
21 I had the first patient who had a tendon rupture. So  
22 before you actually experience a complication, before

1 it's ever happened, you aren't as aware of it in  
2 day-to-day practice, at least I wasn't.

3           Once I knew one patient had a tendon  
4 rupture, I think that you become a little bit more  
5 concerned, you become a little bit more attuned to the  
6 potential for the complication. And not only would  
7 potentially the location of the injection matter, but  
8 also I think one of the patients who had a tendon  
9 rupture was actually on their third injection.

10           There was a patient who had good restoration  
11 of extension, had not yet met that zero to 5 degree  
12 benchmark, but the investigator wanted to try to get  
13 it to that point, did a third injection. The patient  
14 had -- I don't remember the -- had a contracture  
15 probably in the 15 to 20 degree range, the tendon  
16 rupture occurred. That in my experience, after the  
17 tendon rupture happened, once I didn't have patients  
18 who didn't have a full correction but I couldn't feel  
19 a well-defined cord anymore, I didn't give them any  
20 more injections. And I think that's key to the  
21 training.

22           So it's not just where you're giving the

1 injection, but, hey, what we're treating is the cord.  
2 And if you can't feel the cord, you can't access it  
3 safely, don't give the shot.

4 But I think this is an example of the spiral  
5 cord. Again, the finger is pointing out towards the  
6 left, and there is a little blue marker here and a  
7 blue marker here, and I actually put surgical ink  
8 along that cord tissue. And what we can see is nerve  
9 and artery poking out here, the nerve and artery right  
10 here, and this cord coming from underneath to over  
11 top.

12 So it does. This cord tissue as the various  
13 areas of the fascial anatomy come together, they can  
14 wrap around these, but you can see a web space here.  
15 Here's a finger with the web space right in here. So  
16 right at that web, I think you can still get good  
17 access.

18 It is a small needle. Unlike needle  
19 aponeurotomy, you're not passing that needle back and  
20 forth. You're just injecting it right into that cord,  
21 and if a patient -- and you're doing that without a  
22 local anesthetic, so if the patient has a paresthesia,

1     you can stop, redirect.

2                 DR. DELCONTE:   And I do want to remind, we  
3     did not have any nerve or artery injuries in our  
4     series.

5                 DR. O'NEIL:   Are there other questions from  
6     the panel?   Dr. Mazor.

7                 DR. MAZOR:   When you're talking about  
8     monitoring adverse events if this is approved, can you  
9     talk for just a moment about how you will assure that  
10    there aren't misses, that everything is reported and  
11    captured?   Is there any system in place to maximize  
12    that?   Are you worried at all that you will miss  
13    people?

14                DR. DELCONTE:   Dr. Tursi can show you the  
15    targeted pharmacovigilance program that we've put in,  
16    or we propose to put into place that will minimize  
17    that.   I don't think we can ever totally make sure we  
18    won't miss a case.   But go ahead, Jim.

19                DR. TURSI:   I can certainly reiterate  
20    regarding the enhanced safety monitoring program that  
21    we're suggesting for AA4500.   This was from my main  
22    presentation.   Things we'll include without



1 reiterating the entire slide, things will include the  
2 safety hotline. The aggregate safety review will be  
3 performed monthly for the first year. We will be  
4 specifically looking for potential problems, followed  
5 by quarterly reviews Years 2 to 5. And as I said,  
6 there also will be a follow-up questionnaire in the  
7 event of something like a tendon rupture, so we can  
8 track, gather more information and then potentially  
9 adjust our training program or distribution as we need  
10 to based upon those findings, with the ultimate goal,  
11 of course, being to ensure safe and effective use of  
12 AA450.

13 DR. O'NEIL: Next, Dr. Swartz.

14 DR. SWARTZ: My question is, we've already  
15 talked about treatment for surgery after injection,  
16 but how about the reverse? If a patient presents  
17 having had surgery and still has a contracture, are  
18 they still candidates for collagenase injection, and  
19 if so, what are the caveats?

20 DR. DELCONTE: Yes, we've had a number of  
21 patients who have had prior surgery that were entered  
22 into the clinical trial, and we've actually analyzed

1 the data. Here's the response rates. This is overall  
2 in the pooling of the three large or the double-blind  
3 trials, where about 63 percent in patients without  
4 prior surgery, they're in that same range. And then  
5 patients who've had surgery are about 60 percent.

6 And then if we can build this, we further  
7 looked -- because of the way we collected the data, we  
8 also looked at if they had prior surgery in the same  
9 finger. And there's really no overall difference in  
10 patients who had had prior surgery versus patients  
11 with no surgery.

12 DR. O'NEIL: Someone else had a question  
13 over here. No? Okay.

14 Any other questions? Oh, you, my neighbor,  
15 Dr. Buckley.

16 DR. BUCKLEY: Can you give me a little bit  
17 more detail on the safety monitoring post-marketing?  
18 So I'm looking at the slide and it'll be a safety  
19 hotline, which I assume would be for both physicians  
20 and patients to call in, and aggregate safety review  
21 monthly and then quarterly. Are these going to be  
22 questionnaires directly to the physicians who did the

1 procedures?

2           Are patients going to be surveyed? I'd  
3 imagine if patients had a procedure, they might not be  
4 coming back for regular follow-up a year, two years or  
5 five years later. So how do you get that data other  
6 than patients remembering to call in or remembering  
7 that there is a hotline? Will there be some kind of  
8 regular survey both in terms of results and in terms  
9 of adverse events?

10           DR. DELCONTE: There wasn't a regular survey  
11 for patients.

12           Jim, do you want to address that?

13           What we plan to do in the targeted  
14 pharmacovigilance is part of the patient information  
15 brochure. We'll actually indicate what some of the  
16 side effects are to look for, and then we will have a  
17 hotline which will be available for physicians as well  
18 as patients. And we'll be able to transfer these  
19 directly to our safety group for evaluation, whether  
20 it's patients or physicians.

21           If patients aren't following the  
22 instructions and don't return, we don't have a

1 mechanism for that. It's the ones that do return that  
2 we have the mechanism for.

3 DR. BUCKLEY: So -- but am I wrong in  
4 thinking that many patients might not return? It's  
5 not like a rheumatoid arthritis patient who's coming  
6 in every three months for monitoring. If they're  
7 seeing someone for a surgical procedure, unless -- I'm  
8 curious about if you have estimates about how many of  
9 those patients are going to be coming back. Is there  
10 some protocol that you'll be following a certain group  
11 of these patients every three months or once a year,  
12 and how of those patients would it be?

13 DR. T. KAPLAN: I think that -- and I have  
14 another just kind of example, but with collagenase,  
15 all the tendon ruptures happened relatively soon after  
16 the treatment was given, within one to two weeks.

17 Another example that I run into is now with  
18 distal radius fractures, plating of distal radius  
19 fractures, where we put a metallic plate on the  
20 surface of the radius in order to stabilize that  
21 fracture, has a risk of tendon rupture. And we've put  
22 them now on the palm side because we think that's more

1 safe, but patients can still later develop a problem.  
2 And I usually the last day I see a patient after their  
3 fracture, say please call me if you start having any  
4 pain on this side of your wrist, and I've had several  
5 patients come back two years, three years after  
6 treatment who've had irritation.

7           And when I took them to surgery to get their  
8 plate out, could actually see areas of the tendon  
9 where it had been ruptured -- where it had been  
10 thinned.

11           It's well-recognized now that patients won't  
12 come back sometimes until a rupture actually happens.  
13 So I think it's difficult to kind of capture every  
14 patient and to baby-sit them completely, but what we  
15 can do is make sure that patients are aware of what to  
16 look out for, make sure that physicians are aware of  
17 what they need to look out for, and provide mechanisms  
18 for them to contact us if something happens.

19           DR. BUCKLEY: I guess I'm still a little  
20 concerned that there isn't a regular way to follow-up  
21 these patients, not just in terms of adverse events  
22 but to know how long they maintain the benefits. It

1 doesn't sound like we have any way other way other  
2 than patients remembering to call us, or somehow  
3 remembering that a year ago they had some material  
4 that they may not have anymore.

5 DR. DELCONTE: And that's the -- I guess,  
6 limitation of any type of therapy, that sometimes  
7 satisfied patients don't come back. Patients with  
8 problems come back, and that's why we've started the  
9 two- to five-year follow-up study, so that's taking  
10 that large cohort we have in the clinical trials and  
11 following them up through five years to get that  
12 long-term result. So that will add to the knowledge  
13 database of what happens long-term both in terms of  
14 recurrence, progression of disease and safety.

15 DR. O'NEIL: All right. If there is no  
16 further discussion, then we will proceed to the next  
17 session, which is to discuss amid the panel members  
18 the questions to the AAC.

19 The FDA has provided us with three  
20 questions. The first is as follows: Investigator  
21 training in the clinical studies included injection  
22 technique instruction via manuals and DVDs, workshops

1 and investigator meetings. This may be more extensive  
2 than the training proposed for the education of  
3 healthcare professionals in clinical practice if the  
4 product is improved.

5 They ask us to please discuss the adequacy  
6 of the proposed training.

7 And I think an easy -- well, you want to --  
8 I thought an easy way to do this might be to go  
9 around, so we'll start with you, Dr. Weisman.

10 DR. WEISMAN: I think the answer to this  
11 question has to be put in the overall context of what  
12 the mitigation strategies are that we're going to  
13 suggest, and that's how I could answer it. Kathleen  
14 has advised us that the mitigation strategy should be  
15 commensurate with the risk. It shouldn't severely  
16 restrict access, and it shouldn't be burdensome on the  
17 healthcare system.

18 So thinking about this issue and the  
19 discussion around the room and the table, obviously, a  
20 suggestion like restrict this procedure to only  
21 board-certified hand surgeons or certified hand  
22 surgeon, that would be unduly restrictive of access.

1           On the other hand, leaving the whole process  
2   to a voluntarily system that was based upon something  
3   that worked with a highly selective group of skilled  
4   individuals, and then extrapolate that to an  
5   unselected group of individuals, where we don't know  
6   whether it's going to work or not, might be too loose.  
7   So that would be the extremes.

8           And so as I'm thinking about the discussion  
9   here, I'm thinking that what really fits the ideal to  
10  me way to do risk management here would be a mandatory  
11  registry. This would answer Lenore's concerns, which  
12  she's brought up several times, that how are we going  
13  to know whether or not the folks that actually get  
14  this procedure are really monitored long-term, because  
15  there's going to be fallout on either end, the ones  
16  that do well and the ones that do poorly.

17          A mandatory registry also has the advantage  
18  of getting data, which we don't have. It also has the  
19  benefit of casting a kind of accountability to  
20  individuals who both the company, the FDA and to  
21  physicians who participate in this, that they know  
22  they're going into a mandatory registry. And so you



1 don't really get into this lightheartedly. So I would  
2 think that with that level of accountability, then it  
3 might work.

4 I'm sorry if I put the cart before the horse  
5 here in answering this question, because I don't think  
6 that the investigator training in the highly selected  
7 group of individuals that participated in this study  
8 is necessarily the appropriate way to go to unselected  
9 individuals out there in the world. And I don't think  
10 we can fix that, because we don't know anything about  
11 how it works. So getting off that stage, I would move  
12 it more toward the idea of a mandatory registry, which  
13 would have those advantages that I just mentioned.

14 DR. O'NEIL: Yes, sir, Dr. Rosebraugh.

15 DR. ROSEBRAUGH: Yes. This is Curt  
16 Rosebraugh. I just want to probe that answer a little  
17 bit more.

18 So I have to tell you, are you saying a  
19 mandatory registry for every patient that would be  
20 treated with this?

21 DR. WEISMAN: At the outset, yes. A  
22 prospective collection of data on the first year or

1 two or the first numbers that our statisticians would  
2 tell us would be appropriate to know exactly in which  
3 direction we're going.

4 DR. ROSEBRAUGH: Okay.

5 DR. WEISMAN: We can figure that out. It  
6 wouldn't be forever, but it would be for the specific  
7 goals of seeing whether or not the risks of this have  
8 exceeded what our expectations are.

9 DR. ROSEBRAUGH: The reason why I'm asking  
10 is, registries come in two flavors. So we have a lot  
11 of drugs like the TNF drugs where we have registries  
12 that are not part of a REMS. They're a part of a  
13 post-marketing requirement where we say, well, you  
14 know, why don't you register a certain number of folks  
15 and let's follow them for a while and get more data.  
16 Then we have registries that are part of these  
17 Elements to Assure Safe Use. And I just want to make  
18 sure you understand that when we talk about two ends  
19 of the spectrum, we consider that pretty far on one  
20 end of the spectrum.

21 In fact, we very seldom have programs where  
22 we register every patient that gets treatment. It's

1 very extreme for us to do that.

2 DR. WEISMAN: Well, it works in Europe,  
3 where the mandatory registries have given us good data  
4 on the risks of anti-TNF drugs. The registries in the  
5 United States have not given us good data, and we  
6 don't rely on it.

7 DR. ROSEBRAUGH: I appreciate your views. I  
8 just want to make sure everybody understands that is  
9 not something we've routinely done, and it would be  
10 one of the more stricter REMS that we've put in place.

11 DR. O'NEIL: Mr. Brackney?

12 MR. BRACKNEY: Well, from a patient's  
13 standpoint, we've talked about it earlier. You don't  
14 want to do anything that's going to limit access.  
15 Even as an orphan drug with a small population, you  
16 still have to make sure there's doctors out there that  
17 can administer the drug, because clearly, there is an  
18 advantage to people with the disease to have this  
19 treatment as opposed to surgery. So I would be  
20 concerned that the training is sufficient and the base  
21 of the doctors available is as widespread as possible.  
22 But I would at the other end get worried

1    when there's somebody says other.  When I see an other  
2    category on their registry for certification with the  
3    docs, then I worry about how is the other and are  
4    they, back to the point, trainable?  I mean, no  
5    offense, but not every doc is trainable.

6                So I would say sure, beef this up as much as  
7    you can, and then be very selective at the outset  
8    going out with who you have doing it and the doctor  
9    and the practice, and as much as we can, register the  
10   patient so we know what the outcomes are of the people  
11   that are administering the drug, so that we know there  
12   is benefit and that we don't have a hidden problem  
13   somewhere for an untrained person giving -- not taking  
14   the training correctly and administering the drug.

15               DR. O'NEIL:  Dr. Weisman had a reply.

16               DR. WEISMAN:  Can I answer the question?  
17   Now, it's only after years of concern that now we're  
18   getting to the point where Congress is mandating  
19   registries of drug replacements.  And they're very  
20   concerned about outcomes of hip and knee replacements,  
21   and that's being fed back into large grants being  
22   announced by the AHRQ and other organizations after so

1 many years of the voluntary registries not giving us  
2 the information in the United States that we need.

3 And so I wanted just to respond that  
4 voluntary registries have not been very useful  
5 to -- and I understand your concern about perhaps the  
6 onerous issues of having to maintain it, but that  
7 could be a subject of negotiation between yourselves  
8 and the sponsor as to how that actually gets carried  
9 out. But taking it up that level I think is something  
10 that should be considered by the panel here.

11 DR. O'NEIL: Ms. Aronson.

12 MS. ARONSON: I guess I'm trying to weigh in  
13 on the words "may be" in the second sentence. So it's  
14 a little confusing about the -- it doesn't say is  
15 more, will not be as extensive, and I believe that's  
16 the presentation that we had. So I just wanted to be  
17 clear on that "may be." That, for instance, I don't  
18 think there were investigator meetings, and I'm not  
19 sure what other things might be dropped from the list  
20 of training.

21 DR. SAAG: I want to largely second what  
22 Dr. Weisman has said, and I do recognize the FDA's

1 viewpoint on the costs and consequences of  
2 comprehensive registries. But I think as Michael has  
3 illustrated, it's time from a public health  
4 perspective to contemplate new models, whether the  
5 sentinel nodes or some other similar mechanism that is  
6 soon to get started might provide sufficient  
7 surveillance to look at some sample, not a voluntary  
8 registry but some sample of patients who are started  
9 on this therapy, particularly those who might be  
10 treated by physicians with less historic expertise in  
11 doing such procedures, could be done as something the  
12 FDA will have to consider.

13 But that would be what I would consider it  
14 optimal. And I guess it relates to getting back to  
15 the question, and I would answer the question as "no"  
16 in terms of rheumatologists. The average  
17 rheumatologist does not have enough knowledge of the  
18 anatomy of the hand and experience performing  
19 manipulations after injections, or managing,  
20 differentiating postinjection inflammation versus  
21 infection to without significant training be able to  
22 safely administer this product.

1           I think that there is reason to think that  
2   with substantial training that there would  
3   rheumatologists that I would feel confident doing  
4   this, but short of a more extensive training program,  
5   I would have serious reservations about the average  
6   rheumatologist administering this product.

7           DR. O'NEIL: Dr. Buckley.

8           DR. BUCKLEY: I think I'm essentially in  
9   agreement. I think that this study has showed us that  
10  there is a real role for this drug, and I think it's  
11  going to be a very beneficial treatment. But I think  
12  the data that we have is on its use and the results of  
13  its use with hand surgeons and orthopedic surgeons,  
14  and we just don't have the data here to tell us  
15  whether other kinds of physicians, including  
16  rheumatologists, will get these same results.

17           I hope that's true, but I think that if we  
18  sort of jump ahead to the next question, unless  
19  there's data to say that's true, I wouldn't feel  
20  comfortable saying it's a leap of faith, but we think  
21  they can do it based on the number of rheumatologists  
22  in this study.

1 DR. O'NEIL: Dr. Olsen.

2 DR. OLSEN: Well, I have a slightly  
3 different take. I think that the benefits look  
4 significant and the risks look low, and I think that  
5 the plans that have been proposed -- I wasn't -- I  
6 didn't have that opinion before I came to this  
7 meeting, but after having looked at these pictures and  
8 video of demonstrations, I think many of us could be  
9 trained to do this if we felt comfortable doing this.  
10 And we do things every day in our offices that are  
11 totally unregulated. You could put as much  
12 glucocorticoid in as many tendons of somebody's hand  
13 as you wanted to at the moment, and that's probably a  
14 higher risk.

15 So I think by going through what's being  
16 proposed here, registering, have limited access, I  
17 think it sounds like something that would work and  
18 would make something available to a relatively small  
19 number of people who sound like they need it.

20 DR. O'NEIL: And by registering, you mean  
21 registering the healthcare provider who delivers?

22 DR. OLSEN: Oh, I don't want to get into the



1 registry question. I did like the idea of a registry,  
2 but I do understand that's probably a big -- maybe it  
3 could be a sample registry or something like that but  
4 not biased in some way, like figure out some way. The  
5 statisticians could tell us some way to get an  
6 unbiased sample and follow that sample, because I  
7 agree, we need more data. But within the confines of  
8 this being a rare disease and it looking like it has  
9 benefit, I think that shouldn't hold it up.

10 DR. RAPPAPORT: Just to be very clear, the  
11 sample registry is a study, and the other is just  
12 collecting everything and mandating that a patient has  
13 to be registered before they can get it. It's a whole  
14 different ball of wax, but the study is something that  
15 we could, as Dr. Rosebraugh said, do under a  
16 post-marketing requirement.

17 DR. O'NEIL: Dr. McAlindon.

18 DR. McALINDON: So I think when you put this  
19 intervention into the context of the alternative  
20 surgery, the data show that it's relatively safe.  
21 Also, as an orphan drug, I think the primary point of  
22 this is to make it available to people. So I'm

1 concerned about restricting access. There are, of  
2 course, issues of generalizability, but I don't think  
3 necessarily that this panel of hand surgeons is  
4 necessarily generalizable to hand surgeons in the  
5 population.

6           So I think that the training proposed is  
7 likely adequate for clinicians who are accustomed on a  
8 regular basis to doing interventions in the hands.  
9 And I think that some sort of surveillance is  
10 necessary. I think that the registry would be the  
11 gold standard for such surveillance, but an  
12 alternative would be to have the registration happen  
13 at the level of the clinicians so the clinicians would  
14 be registered. And the advantage of that would be  
15 that it would recruit essentially clinicians that had  
16 a more intellectual interest or academic interest in  
17 performing this procedure rather than one in simply  
18 increasing their practice volume.

19           They could then keep a record of patients on  
20 whom they performed this intervention, and that could  
21 be used to address questions which I view as being  
22 perhaps more of a Phase 4 nature, looking at the

1     quality of care and the long-term safety. That to me  
2     would be optimal scenario.

3                 DR. O'NEIL: Dr. Mazor.

4                 DR. MAZOR: I can't talk to the medical and  
5     surgical issues, but I think that one of my concerns  
6     would be that if there is training, and it sounds like  
7     very -- a lot of thought has gone into the training.  
8     My concern would be that people go through the  
9     training and that this issue of kind of doing ones e-  
10    mail simultaneously be somehow addressed, that there  
11    be some sort of check that whatever physician or  
12    surgeon went through it had actually gone through it  
13    and not just signed off on it.

14                And I think what I was trying to ask about  
15    the adverse events before was related to what you're  
16    all calling surveillance and registries, that there  
17    needs -- that these questions really, some of them  
18    aren't answerable at this point. In some way, we need  
19    more data to say, well, is there a difference between  
20    rheumatologists and others in terms of these adverse  
21    outcomes. And I don't know what the options are and  
22    what form that might take, but it seems critical.

1 DR. O'NEIL: Dr. Kaplan.

2 DR. S. KAPLAN: A few thoughts. In terms of  
3 follow-up and monitoring, it's difficult for me to get  
4 patients with problems to come back to the office to  
5 be seen. I don't know how we're going to mandate that  
6 people who are doing well are going to come back. The  
7 Stanford study I referenced was one of the Phase 2  
8 studies. They had 23 people. Nine came back. They  
9 were only able to get nine to come back, one of whom  
10 received placebo. So I don't see how we can easily  
11 monitor this other than keeping in touch with the  
12 providers who do the actual work, to see what kind of  
13 complications they're seeing.

14 As a surgeon, I'm very familiar and  
15 comfortable with credentialing as it relates to  
16 operating-room-based procedures. Delineations of  
17 privileges is something we encounter frequently. The  
18 concept of what we're essentially trying to do here is  
19 credential people to do things in their office.  
20 That's a different world I'm neither familiar nor  
21 comfortable with. There's a lot of things that are  
22 going on in the office -- I agree with Dr. Olsen --

1    that people are doing that we have no idea about and  
2    nobody's watching. People are injecting varicose  
3    veins. There are laser treatments for a variety of  
4    things.

5                I think the onus is on the physician. The  
6    physician states that they're comfortable in this  
7    area, does the appropriate training. I think they are  
8    a licensed physician, they should be credited for  
9    deciding themselves what they're comfortable doing. I  
10   have biases. I think I will do it better than  
11   somebody else. The number three study, Larry Hurst,  
12   he got better results than anybody else. That doesn't  
13   mean other people shouldn't do it.

14               The level of complication, I agree with  
15   Dr. McAlindon. As a surgeon, a tendon rupture is  
16   awful. It's worse than the open procedure for  
17   Dupuytren's, yet at three per 1100, I'm comfortable  
18   with it. It's certainly less common than the rate of  
19   nerve injury either with the needle aponeurotomy or  
20   open surgery. It's less common than the risk of  
21   infection. So I'm comfortable.

22               So this specific question, as I understood

1 the proposal, I think that the training is more than  
2 adequate.

3 DR. O'NEIL: Dr. Swartz.

4 DR. SWARTZ: Thank you. It takes a long  
5 time to see 73 patients with Dupuytren's in most hand  
6 surgeons' practices. Most hand surgeons do five  
7 operations, for the large part: ganglion cyst, carpal  
8 tunnel, trigger finger, de Quervain's releases and  
9 maybe one other procedure. This is a pretty unusual  
10 patient even in a practicing hand surgeon's office.

11 In a rheumatologist's office, in my opinion,  
12 there aren't any patients with rheumatoid arthritis  
13 who have this disease. I've never seen one in 30  
14 years. It's unusual for a rheumatologist to see these  
15 patients. Now, making the diagnosis of a Dupuytren's  
16 nodule instead of a rheumatoid nodule is an important  
17 distinction, but these aren't the patients that will  
18 be treated.

19 So having said that, I think first of all,  
20 the training of video DVD is adequate, that, in my  
21 opinion, the doctors who should take care of these  
22 problems are doctors who see these problems on a

1 regular basis. What the level of their board  
2 certification is is less important than their  
3 familiarity with the disease and its surgical  
4 complications.

5           And lastly, it's my opinion that people who  
6 treat any disease entity should do so if they can  
7 manage the complications of that disease entity. The  
8 complications here are pretty rare, but they're  
9 devastating. A ruptured flexor tendon may not be a  
10 recoverable situation, and a physician bears that  
11 responsibility.

12           So with those caveats, with those warnings  
13 upfront from the company to the doctors they're  
14 marketing to and the information to the patients that  
15 they're going to be providing the medication for, I  
16 think I'm okay with this training and the program  
17 that's been outlined by the company.

18           DR. O'NEIL: Dr. Haque.

19           DR. HAQUE: Thank you. I am also pretty  
20 comfortable with the training regimen that they have,  
21 but I do agree with Dr. Mazor that somehow we have to  
22 enforce that the training's actually done. And I

1 would recommend that the self-assessment exam that the  
2 company has already proposed just be made online, and  
3 that the treating physician actually have to pass it  
4 to get certified. It's the only way that you have of  
5 really enforcing any way that they actually watched  
6 the DVD.

7           As far as healthcare professionals and their  
8 level of training to do this, this actually seems like  
9 a relatively simple procedure. The cords that we're  
10 talking about are usually fairly superficial, as  
11 Dr. Kaplan said, and I think that I don't know how  
12 many rheumatologists actually see patients with  
13 Dupuytren's. I was surprised to hear several  
14 rheumatologists here questioning the ability of the  
15 average rheumatologist to do this procedure, but I  
16 think that this is not going to be such a huge volume  
17 issue that people are going to get rich off of this  
18 procedure.

19           In that situation, I'm more worried about  
20 Dr. Swartz's concern about off-label uses.

21           I think that people who are seeing enough of  
22 this that they actually are willing to take the effort



1 to sign up and get the DVD and take the test are  
2 probably going to be well-qualified to do this.

3 DR. O'NEIL: Dr. Kaplan has another comment.

4 DR. T. KAPLAN: Xiaflex, I guess, comes  
5 under the purview of this rheumatology committee  
6 because, I guess, nobody really knew where to put it.  
7 So the conversation comes up between should it be a  
8 rheumatologist or a hand surgeon.

9 But many of the people I know are in a very  
10 large orthopedic group. They have a hand surgeon or  
11 two. They have a physiatrist or two or three and  
12 maybe a rheumatologist or two, and even  
13 musculoskeletally oriented family practitioners or  
14 internists. I think that's more likely the scenario.  
15 I don't think it's the patient with rheumatoid  
16 arthritis who says to their rheumatologist, oh, by the  
17 way, what is this in my palm? So I think as we think  
18 about it, we think about it as hand surgeons, as  
19 orthopedic surgeons versus people who are caring for  
20 musculoskeletal problems. I think that's a much more  
21 likely scenario, and I'm still comfortable with it.

22 DR. O'NEIL: One comment that I had since I

1   passed my chance on the way through. When you asked  
2   how many rheumatologists actually see these patients,  
3   I'll tell you that as a pediatric rheumatologist, I  
4   haven't seen one since I was a medical student.

5               But the PM&R, the physiatry physicians are  
6   very likely to see some of these patients, I think,  
7   and we should include them in the training program,  
8   because they may be as likely as a rheumatologist,  
9   certainly maybe even more likely.

10              And from my perspective, I think the  
11   proposed training looks very good. After sitting  
12   through this and reading through the information we  
13   were presented with prior to the meeting, I feel like  
14   if I were ever to see one, I might be competent to do  
15   it, having put needles in all kinds of obscene places.

16              So I think that it does look like a fairly  
17   simple procedure. In my mind, I agree completely with  
18   Dr. McAlindon. It looks like it's a very low rate of  
19   although severe complications, it is quite low in good  
20   hands. And hopefully, the training as proposed and  
21   the registry of the trained practitioner will allow  
22   the company to maintain contact with the practitioners

1    who are doing this, and perhaps every few months by e-  
2    mail or by direct mail, inquire of them if they have  
3    seen complications that they need to report, and  
4    thereby sort of enhance reporting of adverse events.

5               We've got a couple more comments from the  
6    docs, and then Dr. Okada had a comment.

7               Dr. Weisman.

8               DR. WEISMAN: To follow up on your comments,  
9    Kathleen, what we've heard is that there's a low rate  
10   of complications, but when it occurs, it's quite  
11   severe, flexor tendon rupture. And if a  
12   rheumatologist does one of those for whatever  
13   procedure they do, if it's injecting an Achilles  
14   tendon sheath, a biceps tendon, it happens once.  
15   They'll really remember that. And the -- so I have  
16   concerns about it.

17              And the other is from our colleagues on the  
18   panel here, they've told us in so many words about the  
19   inadequacy of the follow-up of these patients.  
20   Dr. Kaplan says he doesn't expect to see the ones that  
21   do badly or see the ones that do well, which would be  
22   the majority of people who get the procedure. So the

1 voluntary system of following these patients is really  
2 quite inadequate. There has to be some improvement on  
3 that, just -- that's my response to what I'm hearing  
4 around the table.

5 DR. O'NEIL: Well, I was trying to propose  
6 sort of a middle ground which was enhanced follow-up,  
7 enhanced reporting, which may be working in some other  
8 diseases, but I take your points well, that, yes, we  
9 don't have complete reporting in this country for  
10 virtually anything.

11 Dr. Saag and then Dr. Okada.

12 DR. SAAG: I just want to first of all  
13 clarify my comment from earlier, and I'm not  
14 suggesting that I don't think rheumatologists should  
15 be allowed to do this procedure. But I feel very  
16 strongly that the level of training provided, while  
17 perhaps sufficient for orthopedic surgeons and  
18 particularly for hand surgeons is adequate, I do not  
19 believe at all that this level would be adequate for  
20 most rheumatologists. I would venture to say that  
21 most rheumatologists have no idea where the A-1 pulley  
22 is.

1           And in contrast to comments made about  
2   getting more comfortable as the presentation went on,  
3   I became less comfortable listening to some of the  
4   nuances of how to properly position the injection, and  
5   believe that just watching a DVD and taking a test,  
6   for example, would be fully inadequate in assuring the  
7   appropriate and safe administration of this by  
8   physicians who are not skilled in understanding the  
9   hand anatomy.

10           I think that there are certainly things that  
11   could be done that would not be terribly extensive  
12   that could substantially enhance training, such as  
13   tutorials. There was mention of working with cadavers,  
14   the possibility of developing a model that would  
15   demonstrate the appropriate positioning of the  
16   injections, things that would dramatically improve the  
17   confidence in a physician who normally does not focus  
18   on hand anatomy in administering an injection into the  
19   right location.

20           I would go further to say that most  
21   rheumatologists in practice don't have office staff  
22   that even know how to put on the bulky dressing. Some

1 do work with orthopedic groups. That's true, but many  
2 do not. And there's going to be some training needs  
3 just in understanding the post-procedure care. Again,  
4 I don't believe that a DVD and a examination  
5 afterwards would be sufficient in bringing these  
6 physicians and their office staff up to speed, but I  
7 do think there are things that could be done to  
8 ameliorate that concern.

9 DR. O'NEIL: I'm going to let Dr. Okada go  
10 next, and then we can get to you, Dr. Buckley.

11 DR. OKADA: What I was going to say was  
12 actually touched on by Dr. Rappaport and Dr.  
13 Rosebraugh already. Our concerns related to not  
14 knowing how to generalize the study results are not  
15 ones that necessarily have to be sort of all or  
16 nothing in terms of mandatory registry or nothing. We  
17 do have the post-marketing requirements, and we  
18 could potentially, for example, ask for a large simple  
19 trial, where essentially, you just take all comers of  
20 physicians that would be allowed to utilize the  
21 product, and follow them for a certain period.

22 Something like that might be more feasible

1 and less restrictive on the general public than, say,  
2 a mandatory registry of all patients, so I just wanted  
3 to raise that possibility.

4 DR. O'NEIL: Dr. Buckley.

5 DR. BUCKLEY: I guess I have two other  
6 comments. One is about access and just one is about  
7 the generalizability of this procedure to other types  
8 or a broad variety of providers. I think the -- if  
9 the plan was for this product to have it be a product  
10 that would be used by a broad array of providers, then  
11 I think this study should have been designed to look  
12 at a broad array of providers. As it is, it looked at  
13 very talented array of providers. And I just feel, if  
14 I was designing a treatment that required a certain  
15 level of skill, if I only picked the most skilled  
16 people, then I think I'm going to bias those results  
17 to the best results. That's fine if that's who's  
18 going to be using it.

19 But if really the intent here was this  
20 product was going to be able to be given by many  
21 providers, I think that's the way the trial should  
22 have been designed.

1           And the other thing is, to go back to this  
2   access issue, so if there's a condition that's  
3   prevalent in the population and the provider that you  
4   need to go to, you need to see on a regular basis.  
5   Maybe you just need to see that provider every two or  
6   three months over many, many years. That's a big  
7   access issue if your provider is an hour away, two  
8   hours away or three hours away. If this is a  
9   procedure that will give you a year or many years or a  
10   lifetime benefit for a significant disability, I would  
11   bet in that situation, you'd be more willing to get in  
12   the car and drive an hour and get that procedure done  
13   by somebody who's done it many times.

14           So the access issue, I don't think is quite  
15   the same access issue as, for example, someone with  
16   rheumatoid arthritis or juvenile arthritis who is  
17   really talking about many years of trying to get to a  
18   provider that might be too distant, and I think we  
19   need to weigh that when we think about it.

20           DR. O'NEIL: Dr. Olsen.

21           DR. OLSEN: Well, I just want to point out  
22   that the idea that an initial trial is not broad and



1     that it doesn't include all kinds of scenarios is  
2     exactly what happens in the approval of all medical  
3     interventions that we do. All of the initial TNF  
4     trials excluded people we thought wouldn't get the  
5     drug, and then when the drugs are released, we start  
6     giving to those people and learn new things.

7             I recently did a small trial in  
8     osteoarthritis of a new potential treatment, and I  
9     wanted everyone to do a 25-foot walking time. If you  
10    came in with a walker, I excluded you. Now in real  
11    practice, I'll probably want to see what happens with  
12    those people. But in my first trial, I don't want to  
13    do that. So this is just what you're facing in  
14    trials.

15            So I think that's where a Phase 4 or a  
16    post-marketing trial would be very useful, just  
17    collect more data. It's the same thing that happens  
18    in drugs is what I want to point out.

19            DR. O'NEIL: Dr. Kaplan.

20            DR. SAUL KAPLAN: I do think it is an access  
21    issue. If it requires up to three injections per  
22    joint per affected finger and you can only do one at a

1 time, people don't come in with one affected joint.  
2 They come in with multiple joints, multiple fingers,  
3 both hands. So I think -- and you have to come back  
4 the next day after the injection, so I wouldn't  
5 belittle the access point. I think these are --  
6 multiple visits are going to be involved. More visits  
7 with this procedure potentially than with surgery.

8 DR. O'NEIL: I'd like to ask the  
9 representatives of the -- oh, another comment.

10 DR. McALINDON: Very quickly, access issues  
11 are not necessarily geographic. Insurers can  
12 effectively limit access to quite small domains. If  
13 the one hand surgeon in that domain chooses to not do  
14 this procedure in favor of doing surgery, that could  
15 pose limitation. And so including clinicians who are  
16 perhaps nonsurgical but have some skill in hand  
17 procedures could improve the access.

18 DR. O'NEIL: Now, I'd like to ask the FDA  
19 representatives if there are any other points they  
20 would like for us to address.

21 DR. OKADA: No. Thank you for all that  
22 discussion. That was very helpful.

1                   DR. O'NEIL: All right. Thank you. We will  
2 move on to Question No. 2, which if I can find it  
3 among the many papers I have here, I'll be able to  
4 read to you. This is a voting question, and so I will  
5 first read the question and then give you instructions  
6 regarding voting.

7                   In view of the data available for safety and  
8 efficacy, do you recommend approval of Auxilium's  
9 clostridial collagenase for the treatment of patients  
10 with advanced Dupuytren's disease?

11                  And the voting procedures are as follows:  
12 We will be using the electronic voting system for this  
13 meeting. Each of you have three voting buttons on  
14 your microphone: a yes, a no and an abstain. And  
15 these are flashing before you now. Once we begin the  
16 vote, please press the button that corresponds to your  
17 vote. The vote will then be displayed on the screen.  
18 I will read the vote from the screen into the record.  
19 Next, we will go around the room and each individual  
20 who voted will state their name and the vote into the  
21 record, as well as the reason they voted the way they  
22 did.

1           I will once again read the question, which  
2   you can see on the screen in front of you.

3           In view of the data available for safety and  
4   efficacy, do you recommend approval of Auxilium's  
5   clostridial collagenase for the treatment of patients  
6   with advanced Dupuytren's disease?

7           Please vote.

8           We may have an AV issue.

9   Dr. Haque's -- okay. Good.

10           We're missing one person. So we don't have  
11   a full vote.

12           We will need to repeat the vote. I ask  
13   those who cast their vote to use the identical vote  
14   that they did before. Please don't change your mind  
15   and flip flop, and what we should see now is a  
16   compilation of 12 votes. So if everyone could please  
17   vote now. So when the lights do come on, we will do  
18   the second vote. Please vote.

19           For the record, the voting results are yes,  
20   12; no, zero and abstain, zero to recommend approval.

21           Dr. Haque, would you like to begin stating  
22   your name, your vote and the reason for your vote,

1    please.

2                   DR. HAQUE:   My name is Mustafa Haque, and I  
3   voted yes to approve this medication because I do  
4   think that it will provide significant benefit to  
5   patients, and the overall safety profile looks good.

6                   DR. SWARTZ:   William Swartz, I voted to  
7   approve this drug.   I believe that the risk/benefit  
8   ratio is very low.   The benefit is very high, and I  
9   very much appreciated hearing the testimonials of the  
10  patients that have received this drug.   That did not  
11  necessarily sway my vote, but the vote was made on the  
12  merits of the scientific work presented to us.

13                  DR. S. KAPLAN:   Saul Kaplan, I voted to  
14  approve the use of the drug.   I view it as another  
15  option.   I remain -- or I want to be convinced that  
16  the long-term results are going to hold up enough to  
17  make this something that will become the mainstay of  
18  treatment.   I'm worried that this, like surgery, will  
19  not be the ultimate answer.

20                  DR. MAZOR:   Kathy Mazor, and I voted yes  
21  based on basically the discussion among the physicians  
22  and surgeons, which I again have no medical expertise.

1 The patient testimonials were important about thinking  
2 from the point of a view of a patient. And the  
3 limited understanding I have of the medical  
4 understanding here, it seems like the appropriate  
5 decision and also the FDA's comment that this is also  
6 not a forever decision, that there are additional  
7 studies that could potentially happen in the future  
8 and that things can change if needed.

9 DR. McALINDON: Timothy McAlindon, I voted  
10 yes. There's an acute need for a nonsurgical  
11 intervention for Dupuytren's. This product appears  
12 highly effective, and it has a safety profile that is  
13 acceptable and better than the current surgical  
14 alternative.

15 DR. OLSEN: Nancy Olsen, and I voted yes.  
16 And I agree completely with the comments that were  
17 just made, and I also thought that this satisfied an  
18 unmet need. So it will be very helpful to the  
19 individuals with this disease.

20 DR. BUCKLEY: I'm Lenore Buckley, and I also  
21 voted yes. I think that this is a treatment that  
22 offers patients who have significant disability

1 significant benefits at an acceptable risk.

2 DR. O'NEIL: Kathleen O'Neil, I also voted  
3 yes because this is an effective and reasonably safe  
4 alternative to surgery, and in fact, in some ways may  
5 be better than surgery.

6 DR. SAAG: Ken Saag, I voted yes based on a  
7 highly satisfactory risk/benefit ratio and unmet need.

8 MS. ARONSON: Diane Aronson, I voted yes for  
9 the reasons that have been said.

10 MR. BRACKNEY: Bill Brackney, I voted yes  
11 because it is a better alternative than surgery, and  
12 in the long-term and holds a lot more promise for a  
13 permanent solution than surgery does today.

14 DR. WEISMAN: Michael Weisman, I voted yes  
15 because of the evidence in two very well-done trials  
16 and the significant unmet need.

17 DR. O'NEIL: Thank you, Panel. Now that we  
18 have voted to recommend that this be approved, we are  
19 asked the following questions -- we are asked the  
20 Question 3-A: What additional studies, if any, should  
21 be conducted post-approval to further assess the  
22 safety of the product?

1                   Dr. Weisman, we know you've made up your  
2 mind.

3                   DR. WEISMAN: No.

4                   DR. O'NEIL: No?

5                   DR. WEISMAN: I strongly suggested a  
6 mandatory registry, the details of which can be worked  
7 out as to how what kind of sample and who exactly is  
8 going to do it and pay for it, and how long it needs  
9 to be carried out. I think the statisticians would be  
10 very helpful in that regard. I understand that it's  
11 breaking new ground, as Bob and Curt have told us  
12 since they've really not done this before, and it does  
13 represent at least in their view a somewhat onerous  
14 responsibility.

15                   But on the other hand, what I've tried to  
16 point out is that the voluntary registries that we've  
17 had so far in this country have really been inadequate  
18 to answer the important questions posed by biologic  
19 drugs, even non-steroidal anti-inflammatory drugs and  
20 most all drugs. And, also, the comments from our  
21 colleagues across the table here who've told us about  
22 the routine, usual follow-up of surgical patients or



1 procedure patients is very inadequate. And so that's  
2 the reason I propose this.

3 DR. O'NEIL: Before we proceed with this  
4 portion of the discussion, I asked Nicole to put up  
5 Slide No. 7 first of the FDA's presentation, just to  
6 remind us of the difference between the proposed  
7 post-marketing surveillance that was offered by the  
8 company versus an enforced and mandatory  
9 post-marketing, and these were brought by Dr.  
10 O'Connell.

11 We could -- we are suggested to use some  
12 or -- I'm sorry -- such recommendations may be  
13 important in a setting where one or more of the three  
14 dashed points here are in effect, and I think that the  
15 second dashed point, the product has serious risks  
16 that could affect the patient's decision to use or to  
17 continue to use the product, is applicable to this  
18 particular compound.

19 And then the next slide, just to remind you  
20 that the FDA-approved materials used to aid sponsor  
21 implementation of REMS and/or inform healthcare  
22 providers about serious risks. I'm sorry. The one

1     that I really wanted was the following one.

2                 That we have to remember that mandatory here  
3     is that the FDA requires and enforces this, and then  
4     in Slide 10, that the REMS ETASU program would provide  
5     the most strict control over whether the product is  
6     used per FDA-approved labeling. But the downside is  
7     that it can impose burdens on the healthcare system  
8     and reduce access to care. And so they recommend that  
9     the ETASU program be used only if the product would  
10    otherwise not be approved due to specific serious risk  
11    listed in the labeling.

12                So as we discuss this, we want to make sure  
13    we keep straight what studies need to be done and what  
14    post-marketing should be mandated or used.

15                DR. ROSEBRAUGH: Can you go back a slide?  
16    So let me just kind of go over this a little bit,  
17    because this can be very confusing to people, and I  
18    have to admit it's confusing to me. And so I will  
19    also say that this legislation is sort of a work in  
20    progress, and so we sometimes don't know how to apply  
21    it until we get a case to work on with it.

22                But mandatory enrollment of patients for

1    this particular segment in reality means that in order  
2    for the drug to be used safely, you need to register  
3    the patient and make sure they're followed. So if you  
4    were giving a chronic medicine where you thought it  
5    was vital that you thought they had to have a CBC such  
6    that you would not approve the drug otherwise, then  
7    you would require that patient be enrolled so that we,  
8    we the government, could make sure that they were  
9    getting a monthly CBC. That's really what that means.

10           That's a little bit different than saying we  
11    need more data and I want to know the outcomes of  
12    patients. That is really more a post-marketing study,  
13    where we can say we can require the sponsor to enroll  
14    so many patients in a post-marketing study and say we  
15    want that followed, we want statistical analysis and  
16    all that kind of thing.

17           So these are two different things, and I  
18    just want to make sure people understand it, because  
19    as with any bureaucracy, it can be kind of confusing.

20           DR. O'NEIL: If I might give an example of  
21    mandated follow-up and mandated registry, the  
22    thalidomide story probably fits here as a mandated

1 situation, where physicians are trained in the issues  
2 related to thalidomide. The company will not allow  
3 you to write a prescription without performing that  
4 training. Pharmacists are also registered to dispense  
5 the drug, but only with appropriately trained  
6 physicians -- and particularly in females, pregnancy  
7 tests must be done monthly. And if there is no  
8 evidence of that, the drug cannot be dispensed.

9           So that's a mandated program that's in the  
10 works currently and has been for years.

11           DR. WEISMAN: To try and respond to Curt's  
12 question and I think I understand it, what is our  
13 concern here, the concern really has to do with the  
14 variability of the skills and ability of the  
15 physicians out there to be able to perform this in a  
16 way in which perhaps this voluntary educational  
17 program may not be adequate. We're not sure that  
18 things match. That's, I think, the biggest concern.

19           So what would be the best approach to that  
20 kind of an issue? And I'm not sure that a  
21 post-marketing study really helps us answer that  
22 question. That's where I'm trying to see -- I'm

1     trying to connect the dots here -- or should a  
2     registration situation that you described, where there  
3     is an ability to go back and document and take a look  
4     at what happens to patients going forward might be a  
5     more adequate way of approaching this question.

6             It's not like a situation where we're  
7     looking at risk of a drug or a procedure that's at the  
8     1 percent or below level, where you can survey out  
9     there in a post-marketing situation and where there is  
10    little concern about who's actually giving the drug,  
11    there's more concern about the patient and the  
12    response.

13            Here, there's concern more on the front end,  
14    and that's why I'm bringing this to your attention in  
15    this way. What's the best approach, say -- to ask our  
16    FDA colleagues what would be the best approach that  
17    they think would be most suitable to answer the  
18    question about who is using this drug and what safe  
19    manner, and is the educational approach adequate to  
20    protect us from this? I'm trying to focus this on  
21    what the issue really is.

22            DR. O'NEIL: Dr. Swartz.

1           DR. SWARTZ: I'm not sure that is the issue.  
2   Intellectually, it might be interesting, but the real  
3   issue is what's the real rate of tendon rupture,  
4   because that's the complication. It takes 73 patients  
5   to be treated before one tendon rupture was found in  
6   this study presented by the sponsors. And so it's  
7   going to take a large number of treating physicians to  
8   come up with meaningful numbers over a significant  
9   period of time.

10           And there's another option that I think is  
11   useful than to have a mandated registry, which I think  
12   would be onerous and I'm opposed to. That is, there  
13   are two associations and societies of hand surgeons in  
14   the country that will be taking this on very quickly.  
15   There are academic centers that see large numbers of  
16   patients that will be eager to study these patients  
17   and their treatment thereof. There probably will be  
18   funding dollars provided not only by industry but also  
19   by grants from the societies that are interested in  
20   hand problems, and I think we can get -- while it  
21   won't be the most comprehensive study overall, it'll  
22   be meaningful in what the real rate of tendon rupture

1 is.

2 I think that's a pretty good compromise,  
3 compared with the onerous problem of mandating that  
4 the doctors drag their patients back into the office  
5 over an extended period of time when it's not likely  
6 that that can be done. And there are some precedents  
7 for this sort of thing, and so I would be in favor of  
8 a post-market study that could be done in a hybrid  
9 manner.

10 DR. O'NEIL: But, again, one problem is that  
11 if we do it through the plastics and orthopedic hand  
12 surgery route, we are not going to be capturing the  
13 family practitioner in Elk City, Oklahoma who may have  
14 10 patients in their practice.

15 DR. S. KAPLAN: I bet you will, because the  
16 family practitioner is not going to be repairing the  
17 tendon rupture. So only the farmer who don't want to  
18 take the time to get his tendon rupture repaired will  
19 be lost in that circumstance.

20 DR. O'NEIL: I sit corrected.

21 Dr. Olsen had a comment.

22 DR. RAPPAPORT: Can I make a comment? I

1 think that the concept here is we can do a study, and  
2 we've said this a couple times now. But I just want  
3 to make it clear. We can require a post-marketing  
4 study, and we can talk about what the best way to do  
5 that is, who should be practicing, whether we should  
6 include different specialties and all that versus this  
7 mandatory registry.

8           And in general, we pretty much think that  
9 randomized controlled trials give us better  
10 information, cleaner information about just about  
11 anything. So trying to tease out the type of  
12 information that we'd like to get here about the  
13 safety and who should use this from a registry is  
14 going to be my mind far more difficult than from a  
15 controlled trial.

16           DR. O'NEIL: Dr. Olsen, did you have a  
17 comment?

18           DR. OLSEN: No, I was going to say exactly  
19 that.

20           DR. O'NEIL: Okay. Dr. Buckley.

21           DR. BUCKLEY: I think that if the FDA  
22 decides to approve this drug for use as recommended,



1 then I think a post-marketing study is going to be  
2 necessary. And I think it's going to be necessary to  
3 look at two things. One is safety, and safety across  
4 different kinds of providers, but even within  
5 providers, safety depending on how many injections  
6 that provider does. And also long-term  
7 results, this question, are we going to see rare  
8 systemic allergic reactions that we're really not  
9 going to know about until we get more patients, and  
10 how long are these beneficial results going to last?  
11 And in a real world or in a broader setting, is the  
12 efficacy going to be as good as it looks now?

13 But I think I take a point with the registry  
14 issue, because I think one of the things that this  
15 prospective trial might not tell us is high-risk  
16 groups. What about the person, the high-risk rate of  
17 this in people who have liver disease or alcoholism  
18 patients who might have more of a tendency to clot or  
19 bleed, diabetic patients? I think what these real  
20 world registries can tell us is outside of the defines  
21 of this clinical trial, in the real world, are there  
22 more infections, are there more complications, are

1    there more ruptures if you have diabetics in this  
2    group?

3                   So I think probably the way to go initially  
4    is a post-marketing study.

5                   DR. RAPPAPORT: I actually don't agree,  
6    because you still get into how do you tease out the  
7    background noise from the registry. But as I said, we  
8    can design a trial just about any way we want, and  
9    broaden the enrollment to include people at various  
10   risks. And it'd be a larger trial, but it's going to  
11   give you that information because you got a control in  
12   it. And those are important issues.

13                  DR. SAAG: So I want to put on a  
14   pharmacopoeia head and not take direct issue with what  
15   you're suggesting, Bob, but at least suggest that some  
16   of our current technologies for studying drugs,  
17   devices and biologics maybe are a little bit old-  
18   fashioned. Clinical trials are great for establishing  
19   efficacy, but we know they're terrible for looking at  
20   safety. And when we see a safety signal in a clinical  
21   trial, it should make us particularly concerned about  
22   what's going to happen in the real world.

1           Registries have the limitations of  
2   observational data. What would be ideal here is to do  
3   a large simple comparative effectiveness study. The  
4   problem is there's nothing really to compare. We  
5   don't think that surgery is a good comparator, and I  
6   would be surprised in a Phase 4 study whether you  
7   could really randomize a representative group of  
8   patients. If I saw the results from this and had a  
9   drug that was approved, I wouldn't want to be in a  
10   clinical trial. I'd want to get the real thing.

11           So I think we're in some ways stuck with  
12   some sort of an observational approach, and again,  
13   back to the idea of sentinel nodes, using linked  
14   databases, using large healthcare systems that have  
15   electronic medical records, understanding that there  
16   are issues of confounding by indication and other  
17   things that will be limitations in understanding  
18   safety signals. But I am very concerned about after-  
19   market surveillance and believe that at least at this  
20   point, a registry is going to be necessary, or some  
21   type of a observational design to understand the  
22   safety signal.

1 DR. O'NEIL: Dr. McAlindon.

2 DR. McALINDON: Since it is already proposed  
3 to do some sort of educational intervention with the  
4 clinicians, it would be a fairly simple step to have  
5 all those clinicians registered and have them keep  
6 records on the patients to whom they administer the  
7 intervention with, and undertaking that the patients  
8 will be contacted. They don't necessarily have to be  
9 seen in the office, but they could be contacted by a  
10 mail survey so that we could get more complete data.

11 DR. O'NEIL: Dr. Haque.

12 DR. HAQUE: I agree with Dr. Weisman's point  
13 that the best way to collect the data and really see  
14 what's happening all the way across the board would be  
15 a mandatory registry. But I do think that's a little  
16 bit of an unfair burden on this particular drug when  
17 we don't do it for so many other drugs that also have  
18 very high-risk profiles.

19 And I do think that a broad capture type of  
20 study would be a reasonable way to try to alleviate  
21 some of our concerns that way, although it won't be  
22 perfect. And once again, I would put a plug in for

1   some kind of standardized consent form so patients  
2   really do have an idea of what even to look for,  
3   because, as was mentioned before, if you've spent the  
4   past six years holding your finger down like this and  
5   suddenly you're stuck out here and can't bend it down  
6   again, you may not be unhappy with that, but it  
7   doesn't help us if you don't report it.

8               So patients do need to know really  
9   critically what to look for, and so I do think that  
10  some kind of standardized consent form that informs  
11  them of what their bad outcomes would be would be very  
12  helpful.

13              DR. O'NEIL:   Dr. McAlindon.

14              DR. McALINDON:   I would counsel against  
15  trying to impose a standardized consent form, because  
16  institutions tend to have or view themselves as having  
17  autonomy.  You could promulgate a template, perhaps,  
18  that they could adapt, but I don't see how you could  
19  operate a single design consent form across the  
20  country.

21              DR. O'NEIL:   Dr. Swartz.

22              DR. SWARTZ:   I'll disagree with that.  In

1 plastic surgery circles, we do a certain broad range  
2 of operations but they're pretty standardized,  
3 including breast reductions, tummy tucks, those have -  
4 - the Society of Plastic Surgeons has a very nice  
5 informed consent form. It's not something -- it's  
6 something that it's a tool that you can use, and it's  
7 between the doctor and the patient. The hospital  
8 doesn't have to approve it. You don't even have to  
9 submit it to the hospitals. Most hospitals have their  
10 own individual consent forms that are non-specific.

11 But this way, you have -- you can assure  
12 that the information that's on that sheet is the  
13 information you want imparted, and that is always  
14 between the doctor and the patient to come to an  
15 understanding that they understand that information.  
16 But I think we as clinicians will need -- we can't  
17 manufacture this consent process de novo every time a  
18 patient comes in. So I would urge along with Dr. Haque  
19 that the sponsor provide a patient-friendly, full,  
20 informed consent that we can use.

21 DR. O'NEIL: Dr. Mazor.

22 DR. MAZOR: I just wanted to agree with what

1 Dr. Saag said earlier in terms of the potential value  
2 of using existing databases in some of the large  
3 health plans. It seems like a natural match for this  
4 that might give not a 100 percent of the information  
5 that one would hope for, but an awful lot of it and  
6 would -- a lot of these plans have patients who stay  
7 with them for many, many years, so you would lose some  
8 folks, but you would be able to get some of this  
9 longitudinal data on outcomes that you might not be  
10 expecting at this point. So it seems like something  
11 to consider in post-marketing studies.

12 DR. O'NEIL: Dr. Weisman.

13 DR. WEISMAN: Just to urge some caution  
14 here. I recently saw some data which was very  
15 interesting. It's unpublished but will be soon on  
16 follow-up of patients from a very, very large joint  
17 replacement registry, where they really examine the  
18 question of what are the complications of the patients  
19 that didn't come back to the doctors versus the ones  
20 that did. And it was exactly what you thought, that  
21 the complications were twice as frequent in the  
22 patients that did not come back for follow-up over the

1 same length of time, went to other doctors and so  
2 forth and so on. And this was well-documented.

3 I'm really concerned about the whole system  
4 of voluntary follow-up of these issues. And I think  
5 Ken's point is extremely important here, to understand  
6 that to get good safety data, we're going to need to  
7 be able to apply a very clean mind. As somebody once  
8 told me, the definition of epidemiology is a clean  
9 mind applied to dirty data. So we need to apply a  
10 very clean mind to be able to capture data out there  
11 in those observational cohorts, and I think he's given  
12 you the marching orders about the need to do that.

13 DR. O'NEIL: I think at this point, I'm left  
14 to ask the FDA if they have other questions or other  
15 issues that we have not discussed -- except that I do  
16 have one. I think we need to revisit the question  
17 that Dr. Hamilton rose, that IGE antibody may indeed  
18 be a significant problem as people come back for other  
19 procedures, other injections over time. And we  
20 certainly know that repeated exposure to any foreign  
21 substance, particularly in subcutaneous or mucosal  
22 sites, is going to induce IGE antibody in as efficient



1 way as we know how to do it as humans.

2           So I think we need to address whether it  
3 indeed would be important to either look back through  
4 the sera that may have been collected. I don't know  
5 if it was in the Phase 3 long-term open-label studies,  
6 and also to be very careful about post-marketing  
7 surveillance about allergic and other immunologic  
8 reactions. And I do -- although there has not yet  
9 been a problem with coagulation, I think we need to  
10 keep our mind's eye open to that possibility.

11           DR. DELCONTE: I'd like to ask Paul  
12 Chamberlain to comment, because we have done -- looked  
13 at our serum. We have previously looked at IGE in the  
14 earlier studies. We did not see a correlation, but  
15 Paul.

16           DR. CHAMERLAIN: Yes, thank you for the  
17 question. We really have poured back over the data  
18 from the earlier studies, and the dilemma for us is  
19 that there was no single systemic manifestation of  
20 immediate hypersensitivity even in the re-treated  
21 subjects into open-label studies. So we have no  
22 biological evidence of an IGE-mediated response.

1                   And typically, it's the clinical  
2   manifestation that begins the diagnostic process of  
3   Type 1 hypersensitivity. And if systemic immediate  
4   hypersensitivity reactions were observed, one would do  
5   a skin test, an in vivo test, in favor of an in vitro  
6   test. The in vitro IGE test is very useful perhaps  
7   for a confirmatory analysis where there are clinical  
8   manifestations of potential Type 1 hypersensitivity.

9                   But in the absence of Type 1  
10   hypersensitivity, you have no clinical positive  
11   control for your in vitro IGE analysis. So it's a  
12   little bit of a chicken and egg situation. It's a  
13   dilemma. Without a clinical positive, you've nothing  
14   really to validate the biological sensitivity of your  
15   in vitro analyses. So you can chase a very, very  
16   sensitive bioanalytical method and perhaps pick up  
17   very weak signals, which have no biological relevance  
18   at all.

19                   So this has really been a dilemma for  
20   Auxilium. Can I just refer to the -- actually a  
21   publication from Dr. Hamilton? And Dr. Hamilton did  
22   publish some data in some Peyronie's subjects with an

1 earlier version of AA4500, and of those 45 subjects  
2 tested in a radio binding test, only one out of 44  
3 subjects generated a very, very weak positive in that  
4 assay system. But because the pretreatment sample was  
5 not tested in the same assay, it's impossible to  
6 ascribe that to a treatment-related effect. And  
7 moreover, there were no clinical manifestations in  
8 those subjects.

9           So taking all the data together, the  
10 Auxilium position is that it would not be worthwhile  
11 going back to retrospectively analyze IGE antibodies  
12 in isolation of no clinical manifestation.

13           DR. O'NEIL: So you would propose doing that  
14 only if people had systemic allergic reactions first?

15           DR. DELCONTE: Yes, that's correct.

16           DR. O'NEIL: I would just like to comment,  
17 as someone who did actually complete training in  
18 allergy, that I would be unwilling to let someone do  
19 an intradermal injection in my forearm of clostridium  
20 collagenase.

21           DR. DELCONTE: And we agree that skin  
22 testing is probably not clinically relevant as well.

1           DR. O'NEIL:  So does the FDA have other  
2  issues they would like us to discuss or address?

3           DR. OKADA:  No.  But we would like to once  
4  again express our thanks to the panel for your  
5  participation today, and also for the very helpful  
6  discussion and advice.

7           DR. O'NEIL:  Thank you, everyone.  This  
8  meeting is now adjourned.

9           (Whereupon, at 2:55 p.m., the meeting was  
10 adjourned.)

11

12

13

14

15

16

17

18

19

20

21

22